

A new approach to strong convergence

By CHI-FANG CHEN , JORGE GARZA-VARGAS , JOEL A. TROPP ,
and RAMON VAN HANDEL 

Abstract

A family of random matrices $\mathbf{X}^N = (X_1^N, \dots, X_d^N)$ is said to converge strongly to a family of bounded operators $\mathbf{x} = (x_1, \dots, x_d)$ when $\|P(\mathbf{X}^N, \mathbf{X}^{N*})\| \rightarrow \|P(\mathbf{x}, \mathbf{x}^*)\|$ for every noncommutative polynomial P . This phenomenon plays a key role in several recent breakthroughs on random graphs, geometry, and operator algebras. However, proofs of strong convergence are notoriously delicate and have relied largely on problem-specific methods.

In this paper, we develop a new approach to strong convergence that uses only soft arguments. Our method exploits the fact that for many natural models, the expected trace of $P(\mathbf{X}^N, \mathbf{X}^{N*})$ is a rational function of $\frac{1}{N}$ whose lowest order asymptotics are easily understood. We develop a general technique to deduce strong convergence directly from these inputs using the inequality of A. and V. Markov for univariate polynomials and elementary Fourier analysis. To illustrate the method, we develop the following applications:

1. We give a short proof of the result of Friedman that random regular graphs have a near-optimal spectral gap and obtain a sharp understanding of the large deviations probabilities of the second eigenvalue.
2. We prove a strong quantitative form of the strong convergence property of random permutation matrices due to Bordenave and Collins.
3. We extend the above to any stable representation of the symmetric group, providing many new examples of the strong convergence phenomenon.

1. Introduction

Let $\mathbf{X}^N = (X_1^N, \dots, X_d^N)$ be a sequence of d -tuples of random matrices of increasing dimension, and let $\mathbf{x} = (x_1, \dots, x_d)$ be a d -tuple of bounded

Keywords: Strong convergence, random permutations, random regular graphs
AMS Classification: Primary: 60B20; Secondary: 15B52, 05C80, 46L53, 46L54.
© 2026 Department of Mathematics, Princeton University.

operators. Then \mathbf{X}^N is said to *converge strongly* to \mathbf{x} if

$$\|P(\mathbf{X}^N, \mathbf{X}^{N*})\| \xrightarrow{N \rightarrow \infty} \|P(\mathbf{x}, \mathbf{x}^*)\| \quad \text{in probability}$$

for every noncommutative polynomial P .

The notion of strong convergence has its origin in the work of Haagerup and Thorbjørnsen [31] in the context of complex Gaussian (GUE) matrices. It has since proved to be a powerful tool in a broad range of applications. Notably, strong convergence plays a central role in a series of recent breakthroughs in the study of random lifts of graphs [7], random covering spaces [38], [46], [37], and [48], random minimal surfaces [65], and operator algebras [31], [30], [34], [4], [35], and [8]; we refer to the recent paper [8] for a more extensive discussion and bibliography.

In many cases, norm convergence can be highly nontrivial to establish even for the simplest polynomials P . For example, let $\bar{S}_1^N, \dots, \bar{S}_d^N$ be independent and identically distributed (i.i.d.) random permutation matrices of dimension N . Then the linear polynomial

$$\bar{A}^N = \bar{S}_1^N + \bar{S}_1^{N*} + \dots + \bar{S}_d^N + \bar{S}_d^{N*}$$

is the adjacency matrix of a random $2d$ -regular graph. Every $2d$ -regular graph has largest eigenvalue $2d$ with eigenvector 1_N , while the Alon–Boppana bound states that the second eigenvalue is at least $2\sqrt{2d-1} - o(1)$ [55]. It was conjectured by Alon and proved in a breakthrough monograph of Friedman [27] that the random $2d$ -regular graph satisfies $\|\bar{A}^N|_{\{1_N\}^\perp}\| = 2\sqrt{2d-1} + o(1)$, so that such graphs have near-optimal second eigenvalue. The much more general result that random permutation matrices converge strongly, due to Bordenave and Collins [7], is of major importance in many recent applications cited above.

Given the power of the strong convergence phenomenon, it may be unsurprising that strong convergence has been notoriously difficult to prove (see, e.g., the discussion in [9]). In those cases where strong convergence has been established, the proofs rely on problem-specific methods and often require delicate computations. These obstacles have hampered efforts to establish strong convergence in new situations and to obtain a sharper understanding of this phenomenon.

In this paper, we develop a new approach to strong convergence that uses only soft arguments, and which requires limited problem-specific inputs as compared to previous methods. We will illustrate the method in the context of random permutation matrices and other representations of the symmetric group, resulting in surprisingly short proofs of strong convergence and new quantitative and qualitative information on the strong convergence phenomenon:

- We prove a quantitative form of Friedman’s theorem on the second eigenvalue of random regular graphs [27] that reveals new probabilistic structure. In particular, we obtain a sharp characterization of the tail behavior of the second eigenvalue.
- We obtain a quantitative form of the strong convergence of random permutation matrices due to Bordenave and Collins [7] and [8], which significantly improves the best known convergence rate for arbitrary polynomials.
- We establish strong convergence of random matrices defined by any stable representation of the symmetric group (in the sense of [25]). This provides a large new family of examples of the strong convergence phenomenon, and illustrates that strong convergence arises in settings far beyond the standard representation.

Further applications to several other models will appear in forthcoming work.

Let us emphasize at the outset that the main difficulty in establishing strong convergence is to prove the upper bound

$$\|P(\mathbf{X}^N, \mathbf{X}^{N*})\| \leq \|P(\mathbf{x}, \mathbf{x}^*)\| + o(1) \quad \text{with probability } 1 - o(1)$$

as $N \rightarrow \infty$. The corresponding lower bound then follows automatically, either as a byproduct of the proof or from general principles (see, e.g., [46, pp. 16–19]). We therefore focus on upper bounds throughout this paper, and relegate a brief treatment of the corresponding lower bounds to [Appendix A](#).

Organization of this paper. This paper is organized as follows. In [Section 2](#), we first outline the core ingredients of our approach in a general setting, while [Section 3](#) presents the main results obtained in this paper.

The next two sections introduce some general tools on which our methods are based. [Section 4](#) recalls properties of polynomials and distributions that form the basis for our approach, while [Section 5](#) recalls basic properties of words in random permutation matrices that form the input for our methods.

The rest of the paper is devoted to the proofs of our main results. [Sections 6](#) and [7](#) prove master inequalities for polynomials and smooth functions, respectively. These inequalities are applied in [Section 8](#) to random regular graphs, and in [Section 9](#) to strong convergence of random permutation matrices. [Section 10](#) extends the above results to general stable representations of the symmetric group.

Finally, [Appendix A](#) is devoted to a brief treatment of lower bounds.

Notation. In what follows, $a \lesssim b$ denotes that $a \leq Cb$ for a universal constant C . We denote $\mathbb{N} := \{1, 2, 3, \dots\}$ and $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$.

The symmetric group on N letters is denoted as \mathbf{S}_N , the free group with free generators g_1, \dots, g_d is denoted \mathbf{F}_d , and $e \in \mathbf{F}_d$ denotes the identity element.

We denote by \mathcal{P}_q the space of real polynomials $h : \mathbb{R} \rightarrow \mathbb{R}$ of degree at most q , and by \mathcal{P} the space of all real polynomials. We denote by $h^{(m)}$ the m th derivative of a univariate function, and by $\|h\|_{C^m[a,b]} := \sum_{j=0}^m \sup_{x \in [a,b]} |h^{(j)}(x)|$.

We will denote by $M_N(\mathbb{C})$ the space of $N \times N$ matrices with complex entries. The trace of a matrix M is denoted as $\text{Tr } M$, the normalized trace as $\text{tr}_N M := \frac{1}{N} \text{Tr } M$, and the operator norm as $\|M\|$. The identity matrix is denoted $\mathbf{1}_N \in M_N(\mathbb{C})$, and $\mathbf{1}_N \in \mathbb{C}^N$ denotes the vector all of whose entries equal one. We use the convention that powers bind before the trace, e.g., $\text{tr}_N(X^N)^p := \text{tr}_N[(X^N)^p]$.

If X is a self-adjoint operator and $h : \mathbb{R} \rightarrow \mathbb{C}$, then the operator $h(X)$ is defined by the usual functional calculus (in particular, if X is a self-adjoint matrix, h is applied to the eigenvalues while keeping the eigenvectors fixed).

2. Outline

The main results of this paper will be presented in [Section 3](#), which can be read independently of the present section. However, as the new approach that underlies their proofs is much more broadly applicable, we begin in this section by introducing the core ingredients of the method in a general setting.

Throughout this section, we let X^N be a sequence of N -dimensional self-adjoint random matrices, whose expected empirical eigenvalue distribution converges as $N \rightarrow \infty$ to a limiting probability measure ν_0 with compact support. This *weak convergence* property is captured by convergence of the moments

$$\lim_{N \rightarrow \infty} \mathbf{E}[\text{tr}_N(X^N)^p] = \int x^p \nu_0(dx) \quad \text{for all } p \in \mathbb{N}.$$

The moments $\mathbf{E}[\text{tr}_N(X^N)^p]$ generally admit a combinatorial description whose behavior as $N \rightarrow \infty$ is well understood, so that weak convergence is readily accessible. However, weak convergence can only ensure that a fraction $1 - o(1)$ of eigenvalues converges to the support of ν_0 , and cannot rule out the existence of outliers in the spectrum. This is the basic difficulty in strong convergence problems.

In the problems of interest in this paper, the random matrix $X^N = P(\mathbf{X}^N, \mathbf{X}^{N*})$ is a noncommutative polynomial of a given family

$$\mathbf{X}^N = (X_1^N, \dots, X_d^N)$$

of random matrices. In this setting, it is natural to think of ν_0 itself as the spectral distribution of a certain limiting operator $X_F = P(\mathbf{x}, \mathbf{x}^*)$ in the sense that

$$\tau(X_F^p) = \int x^p \nu_0(dx) \quad \text{for all } p \in \mathbb{N},$$

where $\mathbf{x} = (x_1, \dots, x_d)$ is a limiting family of operators associated to \mathbf{X}^N and τ is a positive linear functional that plays the role of the trace for the limiting operators. Such models are naturally formulated in the setting of C^* -probability spaces, for which we refer to [54] for an excellent introduction. However, this general framework will not be needed in this paper, as the limiting objects associated to the models we will study admit an explicit construction that is recalled in Section 3.1.

Example 2.1. While the aim of this section is to discuss the methods of this paper in a general setting, the reader may keep in mind the following guiding example. Let $X^N = \bar{A}^N|_{\{1_N\}^\perp}$ be the adjacency matrix of a random $2d$ -regular graph with the trivial eigenvalue removed, as defined in the introduction. In this case, the limiting operator X_F may be viewed as the adjacency matrix of the infinite $2d$ -regular tree, and $\tau(X) = \langle \delta_e, X \delta_e \rangle$ where δ_e is the coordinate vector associated with the root of the tree. The spectral distribution ν_0 of X_F is the Kesten–McKay distribution, which is supported in the interval $[-\|X_F\|, \|X_F\|]$ with $\|X_F\| = 2\sqrt{2d-1}$. A more precise description of this model is given in Section 3.2 below.

Remark 2.2. Throughout this paper, we will work with *self-adjoint* random matrices X^N and operators X_F . This entails no loss of generality: since $\|X\|^2 = \|X^*X\|$ for any bounded operator X , strong convergence of arbitrary noncommutative polynomials P is equivalent to strong convergence of the self-adjoint polynomials P^*P . We may therefore restrict attention to self-adjoint polynomials P .

While it is not essential for the applicability of the methods of this paper, we will also assume for simplicity that the random matrices X^N are uniformly bounded, i.e., there is a constant K so that $\|X^N\| \leq K$ almost surely (a.s.) for all N . This is the case for all models considered in this paper; for example, $\|X^N\| \leq 2d$ a.s. in Example 2.1.

In the remainder of this section, we assume that weak convergence

$$(2.1) \quad \lim_{N \rightarrow \infty} \mathbf{E}[\mathrm{tr}_N(X^N)^p] = \tau(X_F^p) \quad \text{for all } p \in \mathbb{N}$$

holds. The problem of *strong convergence* is to show that not only the moments, but also the operator norm, of X^N converges to that of the limiting model X_F , which ensures a lack of outliers in the spectrum. More precisely, in the sequel we aim to explain how to prove the strong convergence upper bound $\|X^N\| \leq \|X_F\| + o(1)$, which is the main difficulty. The corresponding lower bound is typically an easy consequence of (2.1), as will be discussed in Appendix A.

To date, proofs of strong convergence were based on resolvent equations [31] and [64], interpolation [20], [57], [58], and [2], or the moment method [7],

[9], and [8]. The first two approaches rely on analytic tools, such as integration by parts formulae or free stochastic calculus, that are only available for special models. In contrast, the only known way to access the properties of models such as those based on random permutations or group representations is through moment computations. We therefore begin by recalling the challenges faced by the moment method.

2.1. *Prelude: the moment method.* The basic premise behind the moment method is the observation that since $|\tau(X_F^p)| \leq \|X_F\|^p$, (2.1) implies that

$$(2.2) \quad (\mathbf{E}\|X^N\|^{2p})^{1/2p} \leq (\mathbf{E}[\mathrm{Tr}(X^N)^{2p}])^{1/2p} \leq N^{1/2p}(\|X_F\| + o(1))$$

as $N \rightarrow \infty$. This does not in itself suffice to prove $\|X^N\| \leq \|X_F\| + o(1)$, as the right-hand side diverges when p is fixed. The essence of the moment method is that the desired bound would follow if (2.2) remains valid for $p = p(N) \gg \log N$, as then $N^{1/2p} = 1 + o(1)$. We emphasize that this is a much stronger property than (2.1). In practice, the implementation of this method faces two major obstacles.

- While moment asymptotics for fixed p are accessible in many situations, the case where p grows rapidly with N is often a difficult combinatorial problem. Despite the availability of tools to facilitate the analysis of large moments for strong convergence, such as nonbacktracking and linearization methods (see, e.g., [7] and [8]), the core of the analysis remains dependent on delicate computations that do not readily carry over to new situations.
- A more fundamental problem is that the inequality (2.2) that forms the foundation for the moment method may fail to hold altogether for any $p \gg \log N$. To see why, suppose that $\mathbf{P}[\|X^N\| \geq \|X_F\| + \varepsilon] \geq N^{-c}$ for some $\varepsilon, c > 0$. Then

$$\mathbf{E}[\mathrm{Tr}(X^N)^{2p}] \geq N^{-c}(\|X_F\| + \varepsilon)^{2p},$$

which precludes the validity of (2.2) for $p \gg \log N$. It was observed by Friedman [27] that this situation arises already in the setting of random regular graphs due to the presence with probability N^{-c} of dense subgraphs called “tangles”. The application of the moment method in this setting requires conditioning on the absence of tangles, which significantly complicates the analysis.

2.2. *A new approach.* The approach of this paper is also based on moment computations, which are, however, used in an entirely different manner. As we will explain below, our method deduces norm convergence from moments by first letting $N \rightarrow \infty$ and then $p \rightarrow \infty$, bypassing the challenges of the moment method.

Inputs. Our approach requires two basic inputs that we presently describe.

Let $h \in \mathcal{P}_q$ be a polynomial of degree at most q . Then $\mathbf{E}[\mathrm{tr}_N h(X^N)]$ is a linear combination of the moments $\mathbf{E}[\mathrm{tr}_N (X^N)^p]$ for $p \leq q$. We will exploit the fact that in many situations, the moments are rational functions of $\frac{1}{N}$:

$$(2.3) \quad \mathbf{E}[\mathrm{tr}_N h(X^N)] = \Phi_h\left(\frac{1}{N}\right) = \frac{f_h\left(\frac{1}{N}\right)}{g_q\left(\frac{1}{N}\right)},$$

where f_h and g_q are polynomials that depend only on h and q , respectively. This phenomenon is very common; e.g., it arises from Weingarten calculus [21] and [19] or from genus expansions [50, Chapter 1]. For the random permutation models considered in this paper, the relevant properties are recalled in [Section 5](#).

In general, the function Φ_h is extremely complicated. However, our methods will use only soft information on its structure: we require upper bounds on the degrees of f_h and g_q (which are proportional to q for the models considered in this paper), and we must show that g_q does not vanish near zero. Both properties can be read off almost immediately from a combinatorial expression for Φ_h .

The second input to our method is the asymptotic expansion as $N \rightarrow \infty$

$$(2.4) \quad \mathbf{E}[\mathrm{tr}_N h(X^N)] = \nu_0(h) + \frac{\nu_1(h)}{N} + O\left(\frac{1}{N^2}\right),$$

where ν_0 and ν_1 are linear functionals on the space \mathcal{P} of real polynomials. Clearly

$$\begin{aligned} \nu_0(h) &= \Phi_h(0) = \lim_{N \rightarrow \infty} \mathbf{E}[\mathrm{tr}_N h(X^N)] = \tau(h(X_F)), \\ \nu_1(h) &= \Phi'_h(0) = \lim_{N \rightarrow \infty} N(\mathbf{E}[\mathrm{tr}_N h(X^N)] - \tau(h(X_F))) \end{aligned}$$

are defined by the lowest-order asymptotics of the moments. We will exploit that simple combinatorial expressions for ν_0 and ν_1 are readily accessible in practice.

Remark 2.3. It is evident from the above formulas that $\nu_0(h) = \int h(x) \nu_0(dx)$ coincides with the spectral distribution ν_0 of X_F , so we know *a priori* that it is defined by a probability measure. In particular, even though $\nu_0(h)$ appears in (2.4) only for polynomial test functions h , its definition automatically extends to any continuous function h . In contrast, there is no reason *a priori* to expect that $\nu_1(h)$ can be meaningfully defined for non-polynomial test functions h : there could be functions h for which weak convergence holds at a slower rate than $\frac{1}{N}$, in which case the expansion (2.4) fails to hold. That (2.4) and the definition of $\nu_1(h)$ nonetheless extend to *smooth* functions h will be a key output of our method.

Outline. Our basic aim is to achieve the following.

- (a) We aim to show that the validity of (2.4) extends from polynomials to smooth functions h . In particular, we will show that ν_1 extends to a compactly supported distribution (in the sense of Schwartz, cf. Definition 4.6).
- (b) We aim to show that $\text{supp } \nu_1 \subseteq [-\|X_F\|, \|X_F\|]$. (That $\text{supp } \nu_0 \subseteq [-\|X_F\|, \|X_F\|]$ holds automatically as ν_0 is the spectral distribution of X_F .)

A bound on the norm then follows easily. Let $\chi : \mathbb{R} \rightarrow [0, 1]$ be a smooth function so that $\chi = 0$ on $[-\|X_F\| - \frac{\varepsilon}{2}, \|X_F\| + \frac{\varepsilon}{2}]$ and $\chi = 1$ on $[-\|X_F\| - \varepsilon, \|X_F\| + \varepsilon]^c$. Then $\nu_0(\chi) = \nu_1(\chi) = 0$ by (b), and thus (a) yields

$$\mathbf{P}[\|X^N\| \geq \|X_F\| + \varepsilon] \leq \mathbf{E}[\text{Tr } \chi(X^N)] = O\left(\frac{1}{N}\right).$$

As $\varepsilon > 0$ was arbitrary, $\|X^N\| \leq \|X_F\| + o(1)$ in probability.

We now explain the steps that will be used to prove (a) and (b).

Step 1: The polynomial method. At the heart of our approach lies a general method to obtain a quantitative form of (2.4): we will show that

$$(2.5) \quad \left| \mathbf{E}[\text{tr}_N h(X^N)] - \nu_0(h) - \frac{\nu_1(h)}{N} \right| \lesssim \frac{q^8}{N^2} \|h\|_{C^0[-K, K]}$$

for any $h \in \mathcal{P}_q$, where $\|X^N\| \leq K$ a.s. for all N . While achieving such a bound by previous methods would require hard analysis, it arises here from a soft argument: we can “upgrade” an asymptotic expansion (2.4) to a strong nonasymptotic bound (2.5) merely by virtue of the fact that Φ_h is rational.

To this end, observe that the left-hand side of (2.5) is nothing other than the remainder in the first-order Taylor expansion of Φ_h at zero, so that

$$\left| \Phi_h\left(\frac{1}{N}\right) - \Phi_h(0) - \frac{1}{N} \Phi_h'(0) \right| \leq \frac{1}{2N^2} \|\Phi_h''\|_{C^0[0, \frac{1}{N}]}$$

This bound appears useless at first sight, as it relies on the behavior of $\Phi_h(x)$ for $x \notin J := \{\frac{1}{N} : N \in \mathbb{N}\}$ where its spectral interpretation is lost. However, as Φ_h is rational, we can control its behavior by its values on J alone by means of classical inequalities for arbitrary univariate polynomials f of degree q :

- The inequality $\|f'\|_{C^0[-1, 1]} \leq q^2 \|f\|_{C^0[-1, 1]}$ of A. and V. Markov (Lemma 4.1).
- A corollary of the Markov inequality that $\|f\|_{C^0[-1, 1]} \lesssim \sup_{x \in I} |f(x)|$ for any set $I \subset [-1, 1]$ with $O(\frac{1}{q^2})$ spacing between its points (Lemma 4.2).

Applying these inequalities to the numerator and denominator of Φ_h yields (2.5) with minimal effort. We emphasize that the only inputs used here are upper bounds on the degrees of f_h and g_q in (2.3), and that g_q does not vanish near zero.

Step 2: The extension problem. We now aim to extend (2.5) to $h \in C^\infty$. This is not merely a technical issue: while $\mathbf{E}[\mathrm{tr}_N h(X^N)]$ and $\nu_0(h) = \tau(h(X_F))$ are defined for $h \in C^\infty$ by functional calculus, it is unclear that $\nu_1(h)$ can even be meaningfully defined when h is not a polynomial (cf. Remark 2.3).

In order to address these issues, we must replace the degree-dependent bound (2.5) by a bound that depends only on the smoothness of the polynomial h . This is achieved as follows. Rather than applying (2.5) directly to $h \in \mathcal{P}_q$, we expand

$$h(x) = \sum_{j=0}^q a_j T_j(K^{-1}x)$$

in terms of Chebyshev polynomials of the first kind T_j , and apply (2.5) to each T_j individually. As Chebyshev polynomials are uniformly bounded by 1, this replaces $q^8 \|h\|_{C^0[-K,K]}$ by $\sum_{j=1}^q j^8 |a_j| \lesssim \|h\|_{C^9[-K,K]}$ in (2.5), where the latter inequality follows by classical Fourier analysis. We therefore obtain

$$(2.6) \quad \left| \mathbf{E}[\mathrm{tr}_N h(X^N)] - \nu_0(h) - \frac{\nu_1(h)}{N} \right| \lesssim \frac{1}{N^2} \|h\|_{C^9[-K,K]}$$

for every polynomial $h \in \mathcal{P}$. (We will in fact use a stronger inequality of Zygmund, cf. Lemma 4.4, that achieves better rates in our main results.)

Since the inequality (2.6) no longer depends on the degree of the polynomial h , it extends to arbitrary smooth functions h as polynomials are dense in C^∞ . In particular, it ensures that the left-hand side extends uniquely to a compactly supported distribution, so that $\nu_1(h)$ can be defined for any $h \in C^\infty$.

Step 3: The asymptotic moment method. It remains to bound the support of ν_1 . To this end, we make fundamental use of a key property of compactly supported distributions: their support is bounded by the exponential growth rate of their moments (Lemma 4.9). In particular, if we define

$$\rho = \limsup_{p \rightarrow \infty} |\nu_1(x^p)|^{1/p} = \limsup_{p \rightarrow \infty} \lim_{N \rightarrow \infty} \left| N (\mathbf{E}[\mathrm{tr}_N (X^N)^p] - \tau(X_F^p)) \right|^{1/p},$$

then $\mathrm{supp} \nu_1 \subseteq [-\rho, \rho]$. Thus our method ultimately reduces to a form of the moment method, but where we first let $N \rightarrow \infty$ and only then $p \rightarrow \infty$.

In contrast to the moments of the random matrix X^N , the moments of ν_1 turn out to be easy to analyze for the models considered in this paper using a simple idea that is inspired by earlier work of Friedman [26, Lemma 2.4]: even though ν_1 does not have a clear spectral interpretation, its moments can be expressed as a sum of products of matrix elements of powers of X_F . As the sum only has polynomially many terms, the desired bound $\rho \leq \|X_F\|$ follows readily.

2.3. The role of cancellations. A surprising feature of our approach is that tangles, which form a fundamental obstruction to the moment method,

are completely ignored. This seems unexpected, as the method is based on an estimate (2.5) for traces of high degree polynomials which are merely linear combinations of moments. Indeed, as was explained in Section 2.1, the presence of tangles causes the random matrix moments of large degree to be *exponentially* larger than their limiting values, that is, $\mathbf{E}[\mathrm{tr}_N(X^N)^{2p}] \geq e^{cp}\tau(X_F^{2p})$ for $p \gg \log N$. In contrast, (2.5) yields a bound on the difference between linear combinations of the random matrix moments and their limiting values that is only *polynomial* in the degree.

The key feature of (2.5) is that it involves a uniform bound $\|h\|_{C^0[-K,K]}$ on the test function h . It therefore yields no useful information on moments of order $p \gg \log N$, as $\|h\|_{C^0[-K,K]} = K^p$ is exponential in p for $h(x) = x^p$. However, it yields a powerful bound for polynomials h that are uniformly bounded on the interval $[-K, K]$ independently of their degree. The estimate (2.5) therefore reveals a new phenomenon: the effect of tangles on the individual moments cancels out when they are combined to form bounded test functions h . One of the key features of the polynomial method is that it is able to capture these cancellations.

2.4. Related work. The observation that norm bounds follow from the validity of (2.4) for *smooth* functions h and a bound on first-order support $\mathrm{supp} \nu_1$ has been widely used since the works of Haagerup and Thorbjørnsen [31] and Schultz [64] (see also the work of Parraud [58] where higher-order expansions are considered). However, these works rely heavily on analytic and operator-algebraic tools that are not available for the kind of models considered in this paper. What is fundamentally new here is that our method achieves this aim using only simple moment computations, which are much more broadly applicable.

The polynomial method that lies at the heart of our approach is inspired by work in complexity theory [1]. We are aware of little precedent for the use of this method in a random matrix context, beside a distantly related idea that appears in Bourgain and Tzafriri [11, Theorem 2.3]. The first author used a variant of this idea in concurrent work to establish weak convergence of certain random unitary matrices that arise in quantum information theory [14] and [15].

We emphasize that the appearance of Chebyshev polynomials in Step 2 above is unrelated to their connection with nonbacktracking walks that is used, for example, in [27, Lemma 2.3]. Indeed, our Chebyshev polynomials are normalized by the a.s. bound K on $\|X^N\|$ (e.g., $2d$ in Friedman's theorem) as opposed to the support of the limiting spectrum $\|X_F\|$ ($2\sqrt{2d-1}$ in Friedman's theorem).

Finally, we mention the interesting works [6] and [23] that develop a method to bound the spectral radius of certain non-Hermitian matrices using

$N \rightarrow \infty$ moment asymptotics with fixed p , similarly avoiding the challenges of the moment method in that setting. However, the methods developed in these works are unrelated to our approach and are not applicable to the problems considered in this paper.

3. Main results

In this section, we formulate and discuss the main results of this paper. As our primary aim is to achieve strong convergence, we will focus the presentation on upper bounds on the norm as explained in [Section 1](#). Let us note, however, that a byproduct of the analysis will also yield a quantitative form of weak convergence, which is of independent interest; see [Corollary 7.2](#) below.

3.1. Preliminaries. Before we turn to our main results, we must briefly recall some basic facts about random permutation matrices and their limiting model. The following definitions will remain in force throughout this paper.

Definition 3.1. Let $\bar{S}_1^N, \dots, \bar{S}_d^N$ be i.i.d. random permutation matrices of dimension N , and denote by $S_i^N := \bar{S}_i^N|_{\{1_N\}^\perp}$ their restriction to the invariant subspace $\{1_N\}^\perp \subset \mathbb{C}^N$. We will often write $\mathbf{S}^N = (S_1^N, \dots, S_d^N)$ and $\mathbf{S}^{N*} = (S_1^{N*}, \dots, S_d^{N*})$.

Definition 3.2. Let $\mathbf{s} = (s_1, \dots, s_d)$ be defined by $s_i := \lambda(g_i)$, where g_1, \dots, g_d are the free generators of \mathbf{F}_d and $\lambda : \mathbf{F}_d \rightarrow B(l^2(\mathbf{F}_d))$ is the left-regular representation defined by $\lambda(g)\delta_h = \delta_{gh}$. Define the vector state $\tau(a) := \langle \delta_e, a\delta_e \rangle$ on $B(l^2(\mathbf{F}_d))$.

The basic weak convergence property of random permutation matrices, due to Nica [[52](#)] (see [Corollary 5.3](#) for a short proof), states that

$$\lim_{N \rightarrow \infty} \mathbf{E}[\mathrm{tr}_N P(\mathbf{S}^N, \mathbf{S}^{N*})] = \tau(P(\mathbf{s}, \mathbf{s}^*))$$

for every noncommutative polynomial P . This property plays the role of [\(2.1\)](#) in [Section 2](#). The aim of the strong convergence problem is to prove that this convergence holds not only for the trace but also for the norm.

The basic inputs to the methods of this paper, as described in [Section 2.2](#), are well known in the present setting. They will be reviewed in [Section 5](#) below.

Remark 3.3. Even though S_i^N are $(N - 1)$ -dimensional matrices defined on $\{1_N\}^\perp$, we will normalize the trace tr_N by N rather than by $N - 1$ as this leads to cleaner expressions for the rational functions that arise in the proof. This makes no difference to our results, and is mainly done for notational convenience.

3.2. Random regular graphs. Let \bar{A}^N be the adjacency matrix of the random $2d$ -regular graph with N vertices defined in [Section 1](#). Then $A^N :=$

$\bar{A}^N|_{\{1_N\}^\perp}$ is defined by the linear polynomial of random permutation matrices

$$A^N := S_1^N + S_1^{N*} + \cdots + S_d^N + S_d^{N*},$$

and the associated limiting model is

$$A_F := s_1 + s_1^* + \cdots + s_d + s_d^*.$$

Note that A_F is nothing other than the adjacency matrix of the Cayley graph of \mathbf{F}_d generated by g_1, \dots, g_d and their inverses, that is, of the $2d$ -regular tree.

It is a classical fact due to Kesten [43] that $\|A_F\| = 2\sqrt{2d-1}$. That

$$\|A^N\| \leq 2\sqrt{2d-1} + o(1) \quad \text{with probability } 1 - o(1)$$

is due to Friedman [27]. Friedman's proof was simplified by Bordenave [5], and a new proof with an improved convergence rate was given by Huang and Yau [42]. Very recently, the latter approach was further developed in the impressive works of Huang, McKenzie and Yau [40] and [41] to achieve the optimal convergence rate.

3.2.1. An effective Friedman theorem. As a first illustration of the methods of this paper, we will give a short new proof of Friedman's theorem in [Section 8.1](#). A direct implementation of the approach described in [Section 2](#) yields the following:

THEOREM 3.4. *For every $d \geq 2$, $N \geq 1$, and $\varepsilon < 2d - 2\sqrt{2d-1}$, we have*

$$\mathbf{P}[\|A^N\| \geq 2\sqrt{2d-1} + \varepsilon] \lesssim \frac{1}{N} \left(\frac{d \log d}{\varepsilon}\right)^8 \log\left(\frac{2ed}{\varepsilon}\right).$$

[Theorem 3.4](#) implies that when d is fixed as $N \rightarrow \infty$, we have¹

$$\|A^N\| \leq 2\sqrt{2d-1} + O_{\mathbf{P}}\left(\left(\frac{\log N}{N}\right)^{1/8}\right).$$

This rate falls short of the optimal $N^{-2/3}$ rate in Friedman's theorem that was very recently established in [40] and [41]. However, the methods of [42], [40], and [41] rely heavily on the special structure of random regular graphs, and it is unclear at present whether they could be applied to the study of strong convergence. In contrast, the methods of this paper will achieve the same rate for arbitrary polynomials of random permutation matrices; see [Section 3.3](#) below.

The quantitative nature of [Theorem 3.4](#) also enables us to consider what happens when $d, N \rightarrow \infty$ simultaneously. It is an old question of Vu [66, §5] whether $\|A^N\| = (1 + o(1))2\sqrt{2d-1}$ with probability $1 - o(1)$ remains valid when $d, N \rightarrow \infty$ in an arbitrary manner. We can now settle this question for the permutation model of random regular graphs that is considered here (see

¹The notation $Z_N = O_{\mathbf{P}}(z_N)$ denotes that $\{Z_N/z_N\}_{N \geq 1}$ is bounded in probability.

[Remark 3.8](#) below for a brief discussion of other models of random regular graphs).

COROLLARY 3.5. $\|A^N\| = (1 + o(1))2\sqrt{2d-1}$ with probability $1 - o(1)$ whenever $N \rightarrow \infty$ and $d = d(N)$ depends on N in an arbitrary manner.

Proof. That the conclusion holds for $d \geq (\log N)^5$ was proved in [12, §3.2.2]. In the complementary regime $d \leq (\log N)^5$, [Theorem 3.4](#) readily yields the upper bound $\|A^N\| \leq (1+o(1))2\sqrt{2d-1}$ with probability $1-o(1)$, while the corresponding lower bound follows from the Alon–Boppana theorem [55]. \square

3.2.2. The staircase theorem. As was explained in [Section 2](#), the approach of this paper only requires an understanding of the first-order term ν_1 in the $\frac{1}{N}$ -expansion of the moments. However, in the setting of random regular graphs, a detailed understanding of the higher-order terms was achieved by Puder [61] using methods of combinatorial group theory. When such additional information is available, the approach of this paper is readily adapted to achieve stronger results.

The following theorem will be proved in [Section 8.2](#) by taking full advantage of the results of [61]. We emphasize that this is the only part of this paper where we will use asymptotics of expected traces beyond the lowest order.

THEOREM 3.6. *Define*

$$\rho_m := \begin{cases} 2\sqrt{2d-1} & \text{for } 2m-1 \leq \sqrt{2d-1}, \\ 2m-1 + \frac{2d-1}{2m-1} & \text{for } 2m-1 > \sqrt{2d-1}, \end{cases}$$

and let

$$m_* := \left\lfloor \frac{1}{2}(\sqrt{2d-1} + 1) \right\rfloor$$

be the largest integer m so that $2m-1 \leq \sqrt{2d-1}$.

Then for every $d \geq 2$, $m_* \leq m \leq d-1$, and $0 < \varepsilon < \rho_{m+1} - \rho_m$, we have

$$\mathbf{P}[\|A^N\| \geq \rho_m + \varepsilon] \leq \frac{C_d}{N^m} \frac{1}{\varepsilon^{4(m+1)}} \log\left(\frac{2e}{\varepsilon}\right)$$

for all $N \geq 1$, and

$$\mathbf{P}[\|A^N\| \geq \rho_m + \varepsilon] \geq \frac{1 - o(1)}{N^m}$$

as $N \rightarrow \infty$. Here C_d is a constant that depends on d only.

[Theorem 3.6](#) reveals an unusual staircase pattern of the large deviations of $\|A^N\|$, which is illustrated in [Figure 3.1](#). The lower bound, which follows from an elementary argument of Friedman [27, Theorem 2.11], arises due to the presence of tangles: Friedman shows that the presence of a vertex with $m > m_*$ self-loops, which happens with probability $\approx N^{1-m}$, gives rise to an




Figure 3.1. Staircase pattern of the tail probabilities of $\|A^N\|$.

outlier in the spectrum at location $\approx \rho_m$. The fact that our upper bound precisely matches this behavior shows that the tail probabilities of $\|A^N\|$ are completely dominated by these events.

Remark 3.7. Let us highlight two further consequences of [Theorem 3.6](#).

1. [Theorem 3.6](#) shows that $\|A^N\| \leq 2\sqrt{2d-1} + \varepsilon$ with probability at least $1 - \frac{c}{N^{m_*}}$ for any $\varepsilon > 0$, closing the gap between the upper and lower bounds in the main result of Friedman’s original monograph [[27](#), p. 2]. This was previously achieved by Friedman and Kohler [[29](#)] by a refinement of Friedman’s methods.
2. The lower bound of [Theorem 3.6](#) shows that the $O(\frac{1}{N})$ probability bound of [Theorem 3.4](#) cannot be improved in general for fixed $\varepsilon > 0$, as this bound is sharp when $m_* = 1$ (that is, when $2 \leq d \leq 4$).

Remark 3.8. The random regular graph model considered in this paper is known as the *permutation model*. Several other models of random regular graphs are also considered in the literature. All these models are known to be contiguous to the permutation model as $N \rightarrow \infty$ for fixed degree $2d$, so that any statement that holds with probability $1 - o(1)$ for the permutation model remains valid with probability $1 - o(1)$ for the other models; cf. [[27](#), pp. 2–3] and the references therein.

It should be emphasized, however, that low probability events are not preserved by contiguity. In particular, the detailed quantitative picture in [Theorem 3.6](#) is specific to the permutation model. The corresponding picture for other models of random regular graphs must be investigated on a case by case basis.

Similarly, the different models of random regular graphs are no longer contiguous when the degree diverges $d \rightarrow \infty$, so that high degree results as in [Corollary 3.5](#) must be investigated separately for each model. Partial results on this problem for the uniform model of random regular graphs may be found in [[3](#)], [[63](#)], and [[36](#)].

3.3. *Strong convergence of random permutation matrices.* The adjacency matrix of a random regular graph is one very special example of a polynomial of random permutation matrices. The much more general fact that the norm of *every* noncommutative polynomial of random permutation matrices converges to that of its limiting model is an important result due to Bordenave and Collins [7] and [8]. Here we give a short proof that yields an effective form of this result.

We will formulate our strong convergence results for general noncommutative polynomials $P \in M_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ with matrix coefficients, that is,

$$P(\mathbf{s}, \mathbf{s}^*) = \sum_w A_w \otimes w(\mathbf{s}, \mathbf{s}^*),$$

where the sum is over a finite set of words w in the symbols $s_1, \dots, s_d, s_1^*, \dots, s_d^*$ and $A_w \in M_D(\mathbb{C})$ are matrix coefficients. The case of scalar coefficients is recovered for $D = 1$, but the more general setting considered here arises often in applications (e.g., in the study of random lifts [7]). For any such polynomial, we define the norm

$$\|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)} = \sup_{\mathbf{U}} \|P(\mathbf{U}, \mathbf{U}^*)\|,$$

where the supremum is taken over all d -tuples of unitary matrices of any dimension. The notation used here agrees with the standard definition in terms of the norm in the full C^* -algebra of the free group; see, e.g., [17, Theorem 7]. However, this is not important for our purposes, and the reader may simply view the above as the definition of the norm. We emphasize that $\|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$ is not the same as $\|P(\mathbf{s}, \mathbf{s}^*)\|$, which corresponds to the reduced C^* -algebra of the free group. For example, for $P(\mathbf{s}, \mathbf{s}^*) = s_1 + s_1^* + \dots + s_d + s_d^*$, we have $\|P(\mathbf{s}, \mathbf{s}^*)\| = 2\sqrt{2d-1}$ and $\|P\|_{C^*(\mathbf{F}_d)} = 2d$. In practice, $\|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$ may be bounded trivially by the sum of the norms of the matrix coefficients of P .

We now state our main result on strong convergence of random permutations.

THEOREM 3.9. *Let $d \geq 2$, and let $P \in M_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ be any self-adjoint noncommutative polynomial of degree q_0 . Then we have*

$$\mathbf{P}[\|P(\mathbf{S}^N, \mathbf{S}^{N*})\| \geq \|P(\mathbf{s}, \mathbf{s}^*)\| + \varepsilon] \lesssim \frac{D}{N} \left(\frac{K q_0 \log d}{\varepsilon} \right)^8 \log \left(\frac{eK}{\varepsilon} \right)$$

for all $\varepsilon < K - \|P(\mathbf{s}, \mathbf{s}^*)\|$, where $K = \|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$.

This theorem will be proved in [Section 9](#). Note that the limiting norm $\|P(\mathbf{s}, \mathbf{s}^*)\|$ can in principle be computed explicitly using the results of [\[44\]](#).

Remark 3.10. The assumption that P is self-adjoint entails no loss of generality: the analogous bound for a non-self-adjoint polynomial P of degree q_0 follows by applying this result to the self-adjoint polynomial P^*P of degree $2q_0$ (cf. [Remark 2.2](#)).

[Theorem 3.9](#) shows that $\|P(\mathbf{S}^N, \mathbf{S}^{N*})\| \leq \|P(\mathbf{s}, \mathbf{s}^*)\| + O_P\left(\left(\frac{\log N}{N}\right)^{1/8}\right)$ for any polynomial P . This significantly improves the best known bound to date, due to Bordenave and Collins [[8](#), [Theorem 1.4](#)], which yields fluctuations of order $\frac{\log \log N}{\log N}$. Our bound can be directly substituted in applications, such as to random lifts of graphs [[7](#), §1.5], to improve the best known convergence rates.

Let us note that for fixed $\varepsilon > 0$, the tail probability of order $\frac{1}{N}$ in [Theorem 3.9](#) cannot be improved in general, as is illustrated by [Theorem 3.6](#) with $d = 2$.

Remark 3.11. While [Theorem 3.9](#) yields much stronger quantitative bounds for fixed D than prior results, it can only be applied when $D = o(N)$. In contrast, it was recently shown in [[8](#), [Corollary 1.5](#)] that strong convergence remains valid in the present setting even for D as large as $N^{(\log N)^{1/2}}$. Such bounds could be achieved using our methods if the supports of the higher-order terms in the $\frac{1}{N}$ -expansion of the moments are still bounded by $\|P(\mathbf{s}, \mathbf{s}^*)\|$. While this is the case for continuous models such as GUE [[58](#)], it is simply false in the present setting: the proof of [Theorem 3.6](#) shows that tangles can already arise in the second-order term. It is an interesting question whether the approach of this paper can be combined with conditioning on the absence of tangles to achieve improved bounds.

3.4. Stable representations of the symmetric group. Let $\pi_N : \mathbf{S}_N \rightarrow B(V_N)$ be a finite-dimensional unitary representation of the symmetric group \mathbf{S}_N . Then we can define random matrices Π_1^N, \dots, Π_d^N of dimension $\dim V_N$ as

$$(3.1) \quad \Pi_i^N := \pi_N(\sigma_i),$$

where $\sigma_1, \dots, \sigma_d$ are i.i.d. uniformly distributed elements of \mathbf{S}_N . When π_N is the standard representation, we recover the random permutation matrices $\Pi_i^N = S_i^N$ as a special case. Other representations, however, capture a much larger family of random matrix models. We will prove strong convergence for random matrices defined by any *stable* representation of \mathbf{S}_N (see [Section 3.4.1](#)), which yields a far-reaching generalization of [Theorem 3.9](#). This result is of interest for several reasons:

- It provides many new examples of the strong convergence phenomenon.
- It shows that strong convergence can be achieved with much less randomness than in the standard representation: Π_i^N uses the same number of

random bits as S_i^N , but has much larger dimension ($\dim V_N \asymp N^\alpha$ with α arbitrarily large). See [9] for analogous questions in the context of the unitary group.

- It provides new evidence in support of long-standing questions on the expansion of random Cayley graphs of the symmetric group, for which extensive numerical evidence and conjectures are available; see [62] and the references therein.
- Random matrices defined by representations other than the standard representation arise in various applications [28] and [33].

Remark 3.12. The question whether strong asymptotic freeness can be derandomized has been investigated in the theoretical computer science literature [51] and [56] by means of pseudorandom permutation matrices. The results of the present section suggest that high-dimensional representations of \mathbf{S}_N can provide a different perspective on such questions. We omit a detailed discussion of the number of random bits needed by our bounds, as significantly stronger results in this direction were subsequently obtained by Cassidy [13] by combining the methods of this paper with new group-theoretic ideas; see Remark 3.16 below.

3.4.1. *Stable representations.* The approach of this paper requires that the moments of the random matrices of interest are rational functions of $\frac{1}{N}$. For this to be the case, we cannot choose an arbitrary representation π_N of \mathbf{S}_N for each N . Instead, we will work with stable representations [25] and [18] that are defined consistently for different N . We briefly introduce the relevant notions here.

Following [32], denote by $\xi_i(\sigma)$ the number of fixed points of σ^i for $\sigma \in \mathbf{S}_N$. The sequence $\xi_1(\sigma), \xi_2(\sigma), \dots$ determines the conjugacy class of σ . If $\pi_N : \mathbf{S}_N \rightarrow B(V_N)$ is any finite-dimensional unitary representation, its character $\sigma \mapsto \text{Tr } \pi_N(\sigma)$ is a class function and can therefore be expressed as a polynomial of $\xi_1(\sigma), \xi_2(\sigma), \dots$

Definition 3.13. A finite-dimensional unitary representation $\pi_N : \mathbf{S}_N \rightarrow B(V_N)$ of \mathbf{S}_N , defined for each $N \geq N_0$, is *stable* if there exists $r \in \mathbb{N}$ and a polynomial $\varphi \in \mathbb{C}[x_1, \dots, x_r]$ so that $\text{Tr } \pi_N(\sigma) = \varphi(\xi_1(\sigma), \dots, \xi_r(\sigma))$ for all $N \geq N_0, \sigma \in \mathbf{S}_N$.

Thus stable representations are representations of \mathbf{S}_N whose characters are defined by the same polynomial φ for all $N \geq N_0$. For example, the standard representation is stable as it satisfies $\text{Tr } \pi_N = \xi_1 - 1$ for all N .

The irreducible stable representations of \mathbf{S}_N can be constructed explicitly as follows. Fix a base partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_\ell > 0) \vdash |\lambda|$ of $|\lambda| = \sum_{i=1}^\ell \lambda_i$.

Then for every $N \geq |\lambda| + \lambda_1$, the irreducible representation of \mathbf{S}_N defined by

$$(3.2) \quad \lambda[N] = (N - |\lambda| \geq \lambda_1 \geq \dots \geq \lambda_\ell) \vdash N$$

is stable. Moreover, it follows from [32, Proposition B.2] that every stable representation in the sense of Definition 3.13 is a direct sum of such irreducible representations defined by fixed base partitions $\lambda^{(1)}, \dots, \lambda^{(s)}$.

3.4.2. Strong convergence. Fix a stable representation $\pi_N: \mathbf{S}_N \rightarrow B(V_N)$ defined for $N \geq N_0$ by a character polynomial $\varphi \in \mathbb{C}[x_1, \dots, x_r]$. We aim to prove strong convergence of the random matrices Π_1^N, \dots, Π_d^N defined by (3.1).

We will not require that π_N is irreducible, but we assume it does not contain the trivial representation. The dimension of Π_i^N is given by

$$D_N := \dim(V_N) = \text{Tr } \pi_N(e) = \varphi(N, N, \dots, N).$$

Thus D_N is a polynomial in N ; we denote its degree by α , so that $D_N \asymp N^\alpha$.

We now formulate our main result on strong convergence of

$$\mathbf{\Pi}^N = (\Pi_1^N, \dots, \Pi_d^N),$$

whose proof is contained in Section 10 below.

THEOREM 3.14. *Let $d \geq 2$, and let $P \in M_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ be any self-adjoint noncommutative polynomial of degree q_0 . Then we have*

$$\mathbf{P}[\|P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})\| \geq \|P(\mathbf{s}, \mathbf{s}^*)\| + \varepsilon] \leq \frac{CD}{N} \left(\frac{Kq_0 \log d}{\varepsilon} \right)^{4(\alpha+1)} \log \left(\frac{eK}{\varepsilon} \right)$$

for all $\varepsilon < K - \|P(\mathbf{s}, \mathbf{s}^*)\|$ and $N \geq N_0$. Here we define $K = \|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$, and C is a constant that depends on the choice of stable representation.

Remark 3.15. It is certainly possible to obtain an explicit expression for C from the proof; we have suppressed the dependence on the choice of stable representation for simplicity of presentation, and we did not optimize this dependence in the proof.

Remark 3.16. The bound $\|P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})\| \leq \|P(\mathbf{s}, \mathbf{s}^*)\| + O_{\mathbf{P}}\left(\left(\frac{\log N}{N}\right)^{1/4(\alpha+1)}\right)$ that follows from Theorem 3.14 becomes weaker when we consider stable representations of increasingly large dimension $D_N \asymp N^\alpha$. This is not a restriction of our method, however, but rather it reflects a gap in the understanding of stable representations at the time that this paper was written [32, Conjecture 1.8]. Important progress in this direction, recently obtained by Cassidy [13], yields an improved probability bound where $\frac{1}{N\varepsilon^{4(\alpha+1)}}$ is replaced by $\frac{1}{N^{\alpha\varepsilon^{8\alpha}}}$, resulting in a convergence rate that is independent of α . As discussed in [13], this yields strong convergence uniformly for all stable representations with $\alpha \leq N^{\frac{1}{20}-\delta}$ for any $\delta > 0$.

Theorem 3.14 can be readily applied to concrete situations. For example, it implies that random $2d$ -regular Schreier graphs defined by the action of \mathbf{S}_N on k -tuples of distinct elements of $\{1, \dots, N\}$ have second eigenvalue $2\sqrt{2d-1} + o(1)$ with probability $1 - o(1)$, settling a question discussed in [32, §1.4]. Indeed, it is not difficult to see (cf. [32, §8]) that the restricted adjacency matrix $A^N = \bar{A}^N|_{\{1^\perp\}}$ of such a random graph can be represented as

$$A^N = \Pi_1^N + \Pi_1^{N*} + \dots + \Pi_d^N + \Pi_d^{N*}$$

for some stable representation of \mathbf{S}_N (which depends on k) that does not contain the trivial representation, so that the conclusion follows immediately as a special case of **Theorem 3.14**. Applications of this model may be found in [28] and [33]. (Let us note for completeness that the case $k = 1$ reduces to Friedman's theorem, while the case $k = 2$ was previously studied by Bordenave and Collins [7, §1.6].)

4. Basic tools

The aim of this section is to introduce the general facts on polynomials and compactly supported distributions that form the basis for the methods of this paper. While most of the tools in this section follow readily from known results, it is useful to state them in the particular form in which they will be needed.

4.1. Markov inequalities. One of the key tools that will be used in our analysis is the following classical inequality of A. Markov and V. Markov. Here we recall that $(2k-1)!! := (2k-1)(2k-3)\dots 5 \cdot 3 \cdot 1 = \frac{(2k)!}{2^k k!}$.

LEMMA 4.1 (Markov inequality). *Let $h \in \mathcal{P}_q$ and $a > 0$, $m \in \mathbb{N}$. Then we have*

$$\|h^{(m)}\|_{C^0[0,a]} \leq \frac{1}{(2m-1)!!} \left(\frac{2q^2}{a}\right)^m \|h\|_{C^0[0,a]}.$$

Proof. Apply [10, p. 256(d)] to $P(x) = h(\frac{a}{2}x + \frac{a}{2})$. □

Two basic issues will arise in our applications of this inequality. First, we will not be able to control the relevant functions on the entire interval $[0, a]$, but only on a discrete subset thereof. This issue will be addressed using a standard interpolation inequality that is itself a direct consequence of the Markov inequality.

LEMMA 4.2 (Interpolation). *Let $h \in \mathcal{P}_q$, and fix a subset $I \subseteq [0, a]$ such that $[0, a] \subseteq I + [-\frac{a}{4q^2}, \frac{a}{4q^2}]$. Then we have*

$$\|h\|_{C^0[0,a]} \leq 2 \sup_{x \in I} |h(x)|.$$

Proof. Apply [16, Lemma 3(i), p. 91] to $P(x) = h(\frac{a}{2}x + \frac{a}{2})$. □

The second issue is that we will aim to apply these inequalities to rational functions rather than to polynomials. However, in the cases of interest to us, the denominator of the rational function will be nearly constant. The following lemma extends the Markov inequality to this setting.

LEMMA 4.3 (Rational Markov). *Let $r = \frac{f}{g}$ be a rational function with $f, g \in \mathcal{P}_q$, and let $a > 0$ and $m \in \mathbb{N}$. Suppose that $c := \sup_{x,y \in [0,a]} \left| \frac{g(x)}{g(y)} \right| < \infty$. Then*

$$\|r^{(m)}\|_{C^0[0,a]} \leq m! \left(\frac{5cq^2}{a} \right)^m \|r\|_{C^0[0,a]}.$$

Proof. Applying the product rule to $f = rg$ yields

$$r^{(m)}g = f^{(m)} - \sum_{k=1}^m \binom{m}{k} r^{(m-k)}g^{(k)}.$$

As $\frac{1}{c} \left\| \frac{h}{g} \right\|_{C^0[0,a]} \leq \frac{\|h\|_{C^0[0,a]}}{\|g\|_{C^0[0,a]}} \leq \left\| \frac{h}{g} \right\|_{C^0[0,a]}$ for every function h , we can estimate

$$\begin{aligned} \|r^{(m)}\|_{C^0[0,a]} &\leq \left\| \frac{f^{(m)}}{g} \right\|_{C^0[0,a]} + \sum_{k=1}^m \binom{m}{k} \|r^{(m-k)}\|_{C^0[0,a]} \left\| \frac{g^{(k)}}{g} \right\|_{C^0[0,a]} \\ &\leq 2c \sum_{k=1}^m \binom{m}{k} \frac{1}{(2k-1)!!} \left(\frac{2q^2}{a} \right)^k \|r^{(m-k)}\|_{C^0[0,a]} \end{aligned}$$

by applying Lemma 4.1 to $f^{(m)}$ and $g^{(k)}$. Here the factor 2 in the second line is due to the fact that the $f^{(m)}$ and $k = m$ terms in the first line yield the same bound.

We now reason by induction. Clearly the conclusion holds $m = 0$. Now suppose the conclusion holds up to $m - 1$. Then the above inequality yields

$$\begin{aligned} \|r^{(m)}\|_{C^0[0,a]} &\leq \left(\frac{5cq^2}{a} \right)^m \|r\|_{C^0[0,a]} \cdot 2c \sum_{k=1}^m \binom{m}{k} \frac{(m-k)!}{(2k-1)!!} \left(\frac{2}{5c} \right)^k \\ &\leq m! \left(\frac{5cq^2}{a} \right)^m \|r\|_{C^0[0,a]} \cdot 2c \left(\cosh \left(\sqrt{\frac{4}{5c}} \right) - 1 \right) \end{aligned}$$

as $\binom{m}{k} \frac{(m-k)!}{(2k-1)!!} = \frac{2^k m!}{(2k)!}$.

Finally, use $c \geq 1$ and $\cosh(\sqrt{x}) - 1 \leq \frac{5x}{8}$ for $x \in [0, 1]$. \square

The above bounds cannot be essentially improved without further assumptions. Lemma 4.1 attains equality for Chebyshev polynomials [10, p. 256(d)]. The optimality of Lemma 4.2 is discussed in [22], while the optimality of Lemma 4.3 is illustrated by considering $r(x) = \frac{1}{u-x}$ on $x \in [0, 1]$ for $u > 1$.

4.2. *Chebyshev polynomials.* Let $h \in \mathcal{P}_q$ and fix $K > 0$. In this section, we will consider the behavior of h on the interval $[-K, K]$.

Denote by T_j the Chebyshev polynomial of the first kind of degree j , defined by $T_j(\cos \theta) = \cos(j\theta)$. Then h can be uniquely expressed as

$$(4.1) \quad h(x) = \sum_{j=0}^q a_j T_j(K^{-1}x)$$

for some real coefficients a_0, \dots, a_q . Note that these coefficients are precisely the Fourier coefficients of the cosine series $h(K \cos \theta)$. We can therefore apply a classical result of Zygmund on absolute convergence of trigonometric series.

LEMMA 4.4 (Zygmund). *Let h be as in (4.1) and define $f(\theta) := h(K \cos \theta)$. Then*

$$|a_0| \leq \|h\|_{C^0[-K, K]},$$

and

$$\sum_{j=1}^q j^m |a_j| \lesssim \beta_* \|f^{(m+1)}\|_{L^\beta[0, 2\pi]}$$

for every $m \in \mathbb{Z}_+$ and $\beta > 1$, where we defined $1/\beta_* := 1 - 1/\beta$.

Proof. That $|a_0| \leq \|h\|_{C^0[-K, K]}$ follows immediately from

$$a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(\theta) d\theta.$$

To obtain the second estimate, we note that

$$f^{(m)}(\theta) = \begin{cases} \sum_{j=0}^q (-1)^{m/2} j^m a_j \cos(j\theta) & \text{for even } m, \\ \sum_{j=0}^q (-1)^{(m+1)/2} j^m a_j \sin(j\theta) & \text{for odd } m. \end{cases}$$

The conclusion follows from [67, p. 242]. □

A direct consequence is the following.

COROLLARY 4.5. *Let h be as in (4.1) and let $m \in \mathbb{Z}_+$. Then*

$$\sum_{j=0}^q j^m |a_j| \leq c_{m, K} \|h\|_{C^{m+1}[-K, K]},$$

where the constant $c_{m, K}$ depends on m, K only.

Proof. That $f(\theta) = h(K \cos \theta)$ satisfies

$$\|f^{(m+1)}\|_{L^\infty[0, 2\pi]} \leq c_{m, K} \|h\|_{C^{m+1}[-K, K]}$$

follows by the chain rule. It remains to apply Lemma 4.4 with $\beta = \infty$. □

4.3. *Compactly supported distributions.* In this paper we will encounter only distributions on \mathbb{R} . We therefore adopt the following definition [39, §2.3].

Definition 4.6. A linear functional ν on $C^\infty(\mathbb{R})$ such that

$$|\nu(f)| \leq c \|f\|_{C^m[-K,K]} \quad \text{for all } f \in C^\infty(\mathbb{R})$$

holds for some $c, K > 0, m \in \mathbb{Z}_+$, is called a *compactly supported distribution*.

The linear functionals that will arise in this paper will not be defined *a priori* on $C^\infty(\mathbb{R})$, but rather only on the space \mathcal{P} of univariate polynomials. It will be therefore essential to understand when linear functionals on \mathcal{P} can be extended to compactly supported distributions. As we have not located the following result in the literature, we prove it here for completeness.

LEMMA 4.7 (Hausdorff moment problem for compactly supported distributions). *Let ν be a linear functional on \mathcal{P} . Then the following are equivalent.*

1. *There exist $c, m, K \geq 0$ so that $|\nu(h)| \leq cq^m \|h\|_{C^0[-K,K]}$ for all $h \in \mathcal{P}_q$, $q \in \mathbb{N}$.*
2. *There exist $c, m, K \geq 0$ so that $|\nu(T_j(K^{-1}\cdot))| \leq cj^m$ for all $j \in \mathbb{N}$.*
3. *The linear functional ν extends uniquely to a compactly supported distribution.*

Only the implication $1 \Rightarrow 3$ will be used in this paper; the equivalence of $2 \Leftrightarrow 3$ appears implicitly in [24, p. 38].

Proof. That $1 \Rightarrow 2$ is immediate as $\|T_j(K^{-1}\cdot)\|_{C^0[-K,K]} = 1$ by the definition of the Chebyshev polynomials T_j .

To prove that $2 \Rightarrow 3$, fix any $h \in \mathcal{P}$ and express it in the form (4.1). Then

$$|\nu(h)| \leq \sum_{j=0}^q |\nu(T_j(K^{-1}\cdot))| |a_j| \leq c \sum_{j=0}^q j^m |a_j| \leq c_{m,K} \|h\|_{C^{m+1}[-K,K]}$$

by Corollary 4.5. Thus the condition of Definition 4.6 is satisfied for all $h \in \mathcal{P}$. As \mathcal{P} is dense in $C^\infty(\mathbb{R})$ with respect to the norm $\|\cdot\|_{C^{m+1}[-K,K]}$, it is clear that ν extends uniquely to a compactly supported distribution.

To prove that $3 \Rightarrow 1$, it suffices by Definition 4.6 to note that $\|h\|_{C^m[-K,K]} \leq c_{m,K} q^{2m} \|h\|_{C^0[-K,K]}$ for every $h \in \mathcal{P}_q$ and $q \in \mathbb{N}$ by Lemma 4.1. \square

Next, we recall the definition of support [39, §2.2].

Definition 4.8. The *support* $\text{supp } \nu$ of a compactly supported distribution ν is the smallest closed set $A \subset \mathbb{R}$ with the property that $\nu(f) = 0$ for every $f \in C^\infty(\mathbb{R})$ such that $f = 0$ in a neighborhood of A .

It is clear that when the condition of Definition 4.6 is satisfied, we must have $\text{supp } \nu \subseteq [-K, K]$. However, the support may in fact be much smaller.

A key property of compactly supported distributions for our purposes is that their support can be bounded by the growth rate of their moments. While the analogous property of measures is straightforward, its validity for distributions requires justification.

LEMMA 4.9. *Let ν be a compactly supported distribution. Then*

$$\text{supp } \nu \subseteq [-\rho, \rho] \quad \text{with} \quad \rho = \limsup_{p \rightarrow \infty} |\nu(x^p)|^{1/p}.$$

Proof. It is clear from the definition of a compactly supported distribution that we must have $\rho < \infty$. Thus the function F defined by

$$F(z) := \sum_{p=0}^{\infty} \frac{\nu(x^p)}{z^{p+1}}$$

is analytic for $z \in \mathbb{C}$ with $|z|$ sufficiently large. It follows from [24, Lemma 1] (or from [64, Theorem 5.4]) that F can be analytically continued to $\mathbb{C} \setminus \text{supp } \nu$ and that $\text{supp } \nu$ is precisely the set of singular points of F . As the above expansion of F is absolutely convergent in a neighborhood of $z = \pm(\rho + \varepsilon)$ for all $\varepsilon > 0$, it follows that $\pm(\rho + \varepsilon) \notin \text{supp } \nu$ for all $\varepsilon > 0$, which yields the conclusion. \square

4.4. *Test functions.* We finally recall a standard construction of smooth approximations of the indicator function $1_{[-\rho, \rho]^c}$ for which the bound of Lemma 4.4 is well controlled; its sharpness is discussed in [39, pp. 19–22]. Recall that $1/\beta_* = 1 - 1/\beta$.

LEMMA 4.10 (Test functions). *Fix $m \in \mathbb{Z}_+$ and $K, \rho, \varepsilon > 0$ so that $\rho + \varepsilon < K$. Then there exists a function $\chi \in C^\infty(\mathbb{R})$ with the following properties:*

1. $\chi(x) \in [0, 1]$ for all x , $\chi(x) = 0$ for $|x| \leq \rho + \frac{\varepsilon}{2}$, and $\chi(x) = 1$ for $|x| \geq \rho + \varepsilon$;
2. $\|f^{(m+1)}\|_{L^\beta[0, 2\pi]} \leq (Cm)^m (\frac{K}{\varepsilon})^{m+1/\beta_*}$ for all $\beta > 1$.

Here $f(\theta) := \chi(K \cos \theta)$, and C is universal constant.

Proof. Let $F \in C^\infty(\mathbb{R})$ be a nonnegative function that is supported in $[0, 1]$ with $\|F\|_{L^1(\mathbb{R})} = 1$. For $a > 0$, define $F_a(x) := \frac{1}{a} F(\frac{x}{a})$ and $H_a(x) := \frac{1}{a} 1_{[0, a]}(x)$, and let

$$h(x) = \int_{-\infty}^x (F_{\delta/2} * \underbrace{H_{\delta/2m} * \cdots * H_{\delta/2m}}_{m \text{ times}})(y) dy.$$

Here $\delta, \varphi > 0$ are parameters that will be chosen below.

Note that $h \in C^\infty(\mathbb{R})$ by construction. Moreover, the integrand in the definition of h is nonnegative, has $L^1(\mathbb{R})$ -norm one, and is supported in $[0, \delta]$. Thus h is nondecreasing, $h(x) = 0$ for $x \leq 0$, and $h(x) = 1$ for $x \geq \delta$.

We define

$$\chi(x) = h\left(\arcsin\left(\frac{x}{K}\right) - \varphi\right) + h\left(-\arcsin\left(\frac{x}{K}\right) - \varphi\right)$$

for $x \in [-K, K]$, and define $\chi(x) = 1$ otherwise. We now choose the parameters $\varphi = \arcsin\left(\frac{\rho+\varepsilon/2}{K}\right)$ and $\delta = \arcsin\left(\frac{\rho+\varepsilon}{K}\right) - \arcsin\left(\frac{\rho+\varepsilon/2}{K}\right)$ so that 1. holds. Moreover, $\chi(x) = 1$ is constant near $x = \pm K$ as $\rho + \varepsilon < K$, so $\chi \in C^\infty(\mathbb{R})$ by the chain rule.

Now note that $f(\theta) = h\left(\frac{\pi}{2} - \theta - \varphi\right) + h\left(\theta - \frac{\pi}{2} - \varphi\right)$ for $\theta \in [0, \pi]$. Therefore

$$\begin{aligned} \|f^{(m+1)}\|_{L^\beta[0, 2\pi]} &\leq 4\|F_{\delta/2} * H'_{\delta/2m} * \cdots * H'_{\delta/2m}\|_{L^\beta(\mathbb{R})} \\ &\leq 4\|F_{\delta/2}\|_{L^\beta(\mathbb{R})} \|H'_{\delta/2m}\|_{L^1(\mathbb{R})}^m \leq (4m)^m \left(\frac{1}{\delta}\right)^{m+1/\beta} 8\|F\|_{L^\infty(\mathbb{R})}, \end{aligned}$$

where the factor 4 in the first line arises by the triangle inequality and by splitting the L^β norm over the intervals $[0, \pi]$ and $[\pi, 2\pi]$, and the second line uses Young's inequality and $H'_a = \frac{1}{a}(\delta_0 - \delta_a)$. Then 2. follows by noting that $\delta \geq \frac{\varepsilon}{2K}$. \square

5. Words in random permutation matrices

The aim of this section is to recall basic facts about random permutations that will form the input to the methods of this paper. These results are implicit in Nica [53], but are derived in simpler and more explicit form by Linial and Puder [45] whose presentation we follow. We emphasize that all the results that are reviewed in this section arise from elementary combinatorial reasoning; we refer the reader to the expository paper [21, §3] for a brief introduction.

We will work with random permutation matrices $\mathbf{S}^N = (S_1^N, \dots, S_d^N)$ and their limiting model $\mathbf{s} = (s_1, \dots, s_d)$ as defined in Section 3.1. In particular, we recall that $s_i := \lambda(g_i)$, where g_1, \dots, g_d are the free generators of \mathbf{F}_d .

It will be convenient to extend these definitions by setting $g_0 := e$, $g_{d+i} := g_i^{-1}$ and analogously $S_0^N := \mathbf{1}$, $S_{d+i}^N := (S_i^N)^{-1} = S_i^{N*}$ and $s_0 := \mathbf{1}$, $s_{d+i} := s_i^{-1} = s_i^*$ for $1 \leq i \leq d$. This convention will be in force in the remainder of the paper.

We aim to understand the expected trace of words in random permutation matrices and their inverses. To formalize this notion, denote by \mathbf{W}_d the set of all finite words in the letters g_0, \dots, g_{2d} . We implicitly identify $w \in \mathbf{W}_d$ with the word map $w: \mathbf{G}^d \rightarrow \mathbf{G}$ it induces on any group \mathbf{G} . Thus $w(g_1, \dots, g_d) \in \mathbf{F}_d$ is the reduction of the word w , while $w(S_1^N, \dots, S_d^N)$ is the random matrix obtained by substituting $g_i \leftarrow S_i^N$ and multiplying these matrices in the order they appear in w .

5.1. *Rationality.* The first property we will need is that the expected trace of any word is a rational function of $\frac{1}{N}$. We emphasize that only very limited

information on this function will be needed, which is collected in the following lemma:

LEMMA 5.1. *Let $w \in \mathbf{W}_d$ be any word of length at most q and $N \geq q$. Then*

$$\mathbf{E}[\mathrm{tr}_N w(\mathbf{S}^N)] = \frac{f_w(\frac{1}{N})}{g_q(\frac{1}{N})},$$

where

$$g_q(x) := (1-x)^{d_1}(1-2x)^{d_2} \cdots (1-(q-1)x)^{d_{q-1}}$$

with $d_j := \min(d, \lfloor \frac{q}{j+1} \rfloor)$, and f_w, g_q are polynomials of degree at most $q(1 + \log d)$.

Proof. Observe that $\mathbf{E}[\mathrm{Tr} w(S_1^N, \dots, S_d^N)]$ is the expected number of fixed points of $w(\bar{S}_1^N, \dots, \bar{S}_d^N)$ minus one (as we restrict to $\{1_N\}^\perp$). Thus [45, eq. (7)] yields

$$\mathbf{E}[\mathrm{tr}_N w(\mathbf{S}^N)] = -\frac{1}{N} + \sum_{\Gamma} \frac{(\frac{1}{N})^{e_{\Gamma}-v_{\Gamma}+1} \prod_{l=1}^{v_{\Gamma}-1} (1 - \frac{l}{N})}{\prod_{j=1}^d \prod_{l=1}^{e_{\Gamma}^j-1} (1 - \frac{l}{N})},$$

where the sum is over a certain collection of connected graphs Γ with $v_{\Gamma} \leq q$ vertices and $e_{\Gamma} \leq q$ edges, and $e_{\Gamma}^j \geq 0$ are integers so that $e_{\Gamma}^1 + \cdots + e_{\Gamma}^d = e_{\Gamma}$ [45, p. 105]. Note that $e_{\Gamma} - v_{\Gamma} + 1 \geq 0$ as Γ is connected.

The denominator inside the sum can be written equivalently as

$$\prod_{j=1}^d \prod_{l=1}^{e_{\Gamma}^j-1} (1 - \frac{l}{N}) = (1 - \frac{1}{N})^{d_{\Gamma}^1} (1 - \frac{2}{N})^{d_{\Gamma}^2} \cdots (1 - \frac{q-1}{N})^{d_{\Gamma}^{q-1}},$$

with $d_i^{\Gamma} := |\{1 \leq j \leq d : e_{\Gamma}^j \geq i+1\}| \leq d_i$ as $e_{\Gamma}^1 + \cdots + e_{\Gamma}^d \leq q$. Thus

$$\begin{aligned} & \mathbf{E}[\mathrm{tr}_N w(\mathbf{S}^N)] \\ &= \frac{-\frac{1}{N} g_q(\frac{1}{N}) + \sum_{\Gamma} (\frac{1}{N})^{e_{\Gamma}-v_{\Gamma}+1} \prod_{l=1}^{v_{\Gamma}-1} (1 - \frac{l}{N}) \prod_{i=1}^{q-1} (1 - \frac{i}{N})^{d_i-d_i^{\Gamma}}}{g_q(\frac{1}{N})}. \end{aligned}$$

To conclude, note that $\sum_{i=1}^{q-1} d_i \leq \int_1^q \min(d, \frac{q}{x}) dx \leq q(1 + \log d) - \min(d, q)$ and $e_{\Gamma} - \sum_{i=1}^{q-1} d_i^{\Gamma} = |\{1 \leq j \leq d : e_{\Gamma}^j \geq 1\}| \leq \min(d, q)$. Thus g_q has degree at most $q(1 + \log d) - 1$ and each term in the sum has degree at most $q(1 + \log d)$. \square

5.2. *First-order asymptotics.* We now recall the first-order asymptotic behavior of expected traces of words in random permutation matrices. The following is a special case of a result of Nica [53]; see [45, p. 124] for a simple proof.

An element of the free group $v \in \mathbf{F}_d$, $v \neq e$ is called a *proper power* if it can be written as $v = w^k$ for some $w \in \mathbf{F}_d$ and $k \geq 2$, and is called a *non-power* otherwise. We denote by $\mathbf{F}_d^{\mathrm{np}}$ the set of non-powers. Every $v \in \mathbf{F}_d$, $v \neq e$ can be written uniquely as $v = w^k$ for some $w \in \mathbf{F}_d^{\mathrm{np}}$ and $k \geq 1$; cf. [47, Prop. I.2.17].

LEMMA 5.2. *Fix a word $w \in \mathbf{W}_d$ that does not reduce to the identity, and express its reduction as $w(g_1, \dots, g_d) = v^k$ with $v \in \mathbf{F}_d^{\text{np}}$ and $k \geq 1$. Then*

$$\lim_{N \rightarrow \infty} N \mathbf{E}[\text{tr}_N w(\mathbf{S}^N)] = \omega(k) - 1,$$

where $\omega(k)$ denotes the number of divisors of k .

Proof. This follows from [53, Corollary 1.3] by noting as in the proof of Lemma 5.1 that $\mathbf{E}[\text{Tr } w(\mathbf{S}^N)]$ is the expected number of fixed points of $w(\bar{\mathbf{S}}^N)$ minus one. \square

In particular, Lemma 5.2 implies the following:

COROLLARY 5.3 (Weak convergence). *For every word $w \in \mathbf{W}_d$, we have*

$$\lim_{N \rightarrow \infty} \mathbf{E}[\text{tr}_N w(\mathbf{S}^N)] = \tau(w(\mathbf{s})).$$

Proof. Lemma 5.2 implies that $\mathbf{E}[\text{tr}_N w(\mathbf{S}^N)] = o(1)$ for any word w that does not reduce to the identity. On the other hand, if w reduces to the identity, then $w(\mathbf{S}^N)$ is the identity matrix on $\{1_N\}^\perp$ and thus $\mathbf{E}[\text{tr}_N w(\mathbf{S}^N)] = 1 - \frac{1}{N}$. The conclusion follows as clearly:

$$\tau(s_{i_1} \cdots s_{i_k}) = \langle \delta_e, \lambda(g_{i_1}) \cdots \lambda(g_{i_k}) \delta_e \rangle = 1_{g_{i_1} \cdots g_{i_k} = e}. \quad \square$$

6. Master inequalities for random permutations I

6.1. *Master inequalities.* In this section, we develop a core ingredient of our method: inequalities for the normalized trace of polynomials of \mathbf{S}^N .

THEOREM 6.1 (Master inequalities). *Fix a self-adjoint noncommutative polynomial $P \in \mathbf{M}_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ of degree q_0 , and let $K = \|P\|_{\mathbf{M}_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$. Then there exists a linear functional ν_i on \mathcal{P} for every $i \in \mathbb{Z}_+$ so that*

$$\left| \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] - \sum_{i=0}^{m-1} \frac{\nu_i(h)}{N^i} \right| \leq \frac{(4qq_0(1 + \log d))^{4m}}{N^m} \|h\|_{C^0[-K, K]}$$

for every $N, m, q \in \mathbb{N}$ and $h \in \mathcal{P}_q$.

An immediate consequence of Theorem 6.1 is the following corollary that we spell out separately for future reference.

COROLLARY 6.2. *In the setting of Theorem 6.1, we have*

$$|\nu_m(h)| \leq (4qq_0(1 + \log d))^{4m} \|h\|_{C^0[-K, K]}$$

for every $m \in \mathbb{Z}_+$, $q \in \mathbb{N}$, and $h \in \mathcal{P}_q$.

While we will need Theorem 6.1 in its full generality in some applications, many strong convergence results will already arise from the case $m = 2$ as was

explained in [Section 2](#). The case $m = 1$ is of independent interest as it provides a quantitative form of [Corollary 5.3](#); see, for example, [Corollary 7.2](#) below.

6.2. *Proof of [Theorem 6.1](#) and [Corollary 6.2](#).* Throughout the proof, we fix a polynomial P as in [Theorem 6.1](#). We emphasize that all the objects that appear in the proof depend implicitly on the choice of P .

We begin by noting that $(\text{tr}_D \otimes \text{id})(h(P(\mathbf{S}^N, \mathbf{S}^{N*})))$ is a linear combination of words $w(\mathbf{S}^N)$ of length at most qq_0 for every $h \in \mathcal{P}_q$. Thus [Lemma 5.1](#) yields

$$\mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] = \Psi_h\left(\frac{1}{N}\right) = \frac{f_h\left(\frac{1}{N}\right)}{g_{qq_0}\left(\frac{1}{N}\right)}$$

for $N \geqq qq_0$, where f_h, g_{qq_0} are polynomials of degree at most $qq_0(1 + \log d)$.

As Ψ_h has no pole at 0, we can define for each $m \in \mathbb{Z}_+$

$$\nu_m(h) := \frac{\Psi_h^{(m)}(0)}{m!}.$$

It is clear from the definition of Ψ_h that ν_m defines a linear functional on \mathcal{P} . Moreover, it follows immediately from Taylor's theorem that

$$(6.1) \quad \left| \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] - \sum_{i=0}^{m-1} \frac{\nu_i(h)}{N^i} \right| \leq \frac{\|\Psi_h^{(m)}\|_{C^0[0, \frac{1}{N}]}}{m!} \frac{1}{N^m},$$

provided N is large enough that Ψ_h has no poles on $[0, \frac{1}{N}]$. The main idea of the proof is that we will use [Lemma 4.3](#) to bound the remainder term in the Taylor expansion. To this end we must show that the function g_{qq_0} is nearly constant.

LEMMA 6.3. $e^{-1} \leqq g_q(x) \leqq 1$ for every $x \in [0, \frac{1}{q^2(1+\log d)}]$ and $q \in \mathbb{N}$.

Proof. The definition of g_q in [Lemma 5.1](#) implies $(1 - qx)^{q(1+\log d)-1} \leqq g_q(x) \leqq 1$ for $x \in [0, \frac{1}{q}]$, where we used that $d_1 + \dots + d_{q-1} \leqq q(1 + \log d) - 1$ as shown in the proof of [Lemma 5.1](#). The conclusion follows using $(1 - \frac{1}{a})^{a-1} \geqq e^{-1}$ for $a > 1$. \square

Next, we must obtain an upper bound on $\|\Psi_h\|_{C^0[0, \frac{1}{N}]}$.

LEMMA 6.4. *The following holds: $\|\Psi_h\|_{C^0[0, \frac{1}{M}]} \leqq 2e\|h\|_{C^0[-K, K]}$ for $M = 2(qq_0(1 + \log d))^2$.*

Proof. As $\|P(\mathbf{S}^N, \mathbf{S}^{N*})\| \leqq \|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$, we have

$$|\Psi_h(\frac{1}{N})| = |\mathbf{E}[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*}))]| \leqq \|h\|_{C^0[-K, K]}$$

for all $N \in \mathbb{N}$. We will extend this bound from the set $I = \{\frac{1}{N} : N \geqq M\}$ to the interval $[0, \frac{1}{M}]$ using polynomial interpolation. First, note that

$$|f_h(\frac{1}{N})| \leqq |\Psi_h(\frac{1}{N})| \leqq \|h\|_{C^0[-K, K]}$$

as $g_{qq_0}(\frac{1}{N}) \leq 1$ for $N \geq q$. The assumption of [Lemma 4.2](#) is satisfied for $h \leftarrow f_h$, $q \leftarrow qq_0(1 + \log d)$, $a = \frac{1}{M}$ as soon as $M \geq 2(qq_0(1 + \log d))^2$, which yields

$$\|f_h\|_{C^0[0, \frac{1}{M}]} \leq 2\|h\|_{C^0[-K, K]}.$$

It remains to apply [Lemma 6.3](#) to estimate $\|\Psi_h\|_{C^0[0, \frac{1}{M}]} \leq e\|f_h\|_{C^0[0, \frac{1}{M}]}$. \square

We can now conclude the proof.

Proof of [Theorem 6.1](#) and [Corollary 6.2](#). Let $M = 2(qq_0(1 + \log d))^2$. Then

$$\frac{\|\Psi_h^{(m)}\|_{C^0[0, \frac{1}{M}]}}{m!} \leq (4qq_0(1 + \log d))^{4m} \|h\|_{C^0[-K, K]}$$

by [Lemmas 4.3](#), [6.3](#), and [6.4](#), where we used that $(10e)^m 2e \leq 4^{4m}$ for $m \geq 1$. Thus the proof of [Theorem 6.1](#) for $N \geq M$ follows directly from [\(6.1\)](#).

The proof of [Corollary 6.2](#) for $m \geq 1$ now follows by multiplying the bound of [Theorem 6.1](#) by N^m and taking $N \rightarrow \infty$. As

$$\nu_0(h) = \lim_N \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right],$$

the conclusion of [Corollary 6.2](#) evidently also holds for $m = 0$.

It remains to prove [Theorem 6.1](#) for $N < M$. To this end, we can trivially bound the left-hand side of the inequality in [Theorem 6.1](#) by

$$\begin{aligned} & \left| \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] - \sum_{i=0}^{m-1} \frac{\nu_i(h)}{N^i} \right| \\ & \leq \left(1 + \sum_{i=0}^{m-1} \frac{(4qq_0(1 + \log d))^{4i}}{N^i} \right) \|h\|_{C^0[-K, K]} \end{aligned}$$

using the triangle inequality and [Corollary 6.2](#). But note that

$$1 < \frac{2^k (qq_0(1 + \log d))^{4k}}{N^k}$$

for every $k \geq 1$ as we assumed $N < M$. We can therefore estimate

$$1 + \sum_{i=0}^{m-1} \frac{(4qq_0(1 + \log d))^{4i}}{N^i} \leq \left(2^m + \sum_{i=0}^{m-1} 4^{4i} 2^{m-i} \right) \frac{(qq_0(1 + \log d))^{4m}}{N^m},$$

and we conclude using $2^m + \sum_{i=0}^{m-1} 4^{4i} 2^{m-i} \leq 4^{4m}$. \square

7. Master inequalities for random permutations II

The aim of this short section is to extend the master inequalities of [Theorem 6.1](#) from polynomials to general smooth test functions. This extension will play a key role in the proofs of strong convergence.

THEOREM 7.1 (Smooth master inequalities). *Fix a self-adjoint noncommutative polynomial $P \in \mathbb{M}_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ of degree q_0 , and let*

$$K = \|P\|_{\mathbb{M}_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}.$$

Then there exists a compactly supported distribution ν_i for every $i \in \mathbb{Z}_+$ so that

$$\left| \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] - \sum_{i=0}^{m-1} \frac{\nu_i(h)}{N^i} \right| \lesssim \frac{(4q_0(1 + \log d))^{4m}}{N^m} \beta_* \|f^{(4m+1)}\|_{L^\beta[0, 2\pi]}$$

for all $N, m \in \mathbb{N}$, $\beta > 1$, $h \in C^\infty(\mathbb{R})$, where $f(\theta) := h(K \cos \theta)$ and $1/\beta_ := 1 - 1/\beta$.*

Proof. We first note that [Corollary 6.2](#) ensures, by [Lemma 4.7](#), that the linear functionals ν_i in [Theorem 6.1](#) extend uniquely to compactly supported distributions.

Let $h \in \mathcal{P}_q$ and express it in the form [\(4.1\)](#). Rather than applying [Theorem 6.1](#) to h directly, we apply it to the Chebyshev polynomials $T_j(K^{-1}\cdot)$ to obtain

$$\left| \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] - \sum_{i=0}^{m-1} \frac{\nu_i(h)}{N^i} \right| \leq \frac{(4q_0(1 + \log d))^{4m}}{N^m} \sum_{j=1}^q j^{4m} |a_j|.$$

Here we used that the left-hand side of [Theorem 6.1](#) vanishes when h is the constant function (as then $\nu_0(h) = h$, $\nu_i(h) = 0$ for $i \geq 1$), so the constant term in the Chebyshev expansion cancels. The proof is completed for $h \in \mathcal{P}_q$ by applying [Lemma 4.4](#). The conclusion extends to $h \in C^\infty$ as \mathcal{P}_q is dense in $C^\infty[-K, K]$. \square

The special case $m = 1$ is of independent interest, as it yields a quantitative formulation of weak convergence (cf. [Corollary 5.3](#)).

COROLLARY 7.2 (Quantitative weak convergence). *In the setting of [Theorem 7.1](#),*

$$\begin{aligned} & \left| \mathbf{E} \left[\text{tr}_{DN} h(P(\mathbf{S}^N, \mathbf{S}^{N*})) \right] - (\text{tr}_D \otimes \tau)(h(P(\mathbf{s}, \mathbf{s}^*)) \right) \right| \\ & \lesssim \frac{(q_0(1 + \log d))^4}{N} \beta_* \|f^{(5)}\|_{L^\beta[0, 2\pi]} \end{aligned}$$

for all $N \in \mathbb{N}$, $\beta > 1$, $h \in C^\infty(\mathbb{R})$, where $f(\theta) := h(K \cos \theta)$ and $1/\beta_ := 1 - 1/\beta$.*

Proof. This is simply a restatement of the special case of [Theorem 7.1](#) with $m = 1$, where we note that $\nu_0(h) = (\text{tr}_D \otimes \tau)(h(P(\mathbf{s}, \mathbf{s}^*)))$ by [Corollary 5.3](#). \square

Remark 7.3. When h is chosen to be a smooth approximation of the indicator function of an interval, [Corollary 7.2](#) shows that the spectral distribution

of $P(\mathbf{S}^N, \mathbf{S}^{N*})$ is well approximated by that of $P(\mathbf{s}, \mathbf{s}^*)$ at a mesoscopic scale. While far sharper results at the local scale are proved in [42], [40], and [41] for random regular graphs, no result of this kind appears to be known to date for arbitrary polynomials P .

In the remainder of this paper, we adopt the notation for the distributions ν_i as in [Theorem 7.1](#). It should be emphasized, however, that the definition of ν_i depends both on the model and on the noncommutative polynomial P under consideration. As each of the following sections will be concerned with a specific model and choice of polynomial P , the meaning of ν_i will always be clear from context.

8. Random regular graphs

The aim of this section is to prove our main results on random regular graphs, the effective Friedman [Theorem 3.4](#) and the staircase [Theorem 3.6](#).

8.1. *Proof of [Theorem 3.4](#).* The proof of [Theorem 3.4](#) is based on [Theorem 7.1](#) with $m = 2$. Let us spell out what this says in the present setting.

LEMMA 8.1. *There exists a compactly supported distribution ν_1 such that for every $N \in \mathbb{N}$, $\beta > 1$, and $h \in C^\infty(\mathbb{R})$, we have*

$$\left| \mathbf{E}[\mathrm{tr}_N h(A^N)] - \tau(h(A_F)) - \frac{1}{N} \nu_1(h) \right| \lesssim \frac{(1 + \log d)^8}{N^2} \beta_* \|f^{(9)}\|_{L^\beta[0, 2\pi]},$$

where $f(\theta) := h(2d \cos \theta)$ and $1/\beta_* = 1 - 1/\beta$.

Proof. Apply [Theorem 7.1](#) with $m = 2$ and $P(\mathbf{s}, \mathbf{s}^*) = s_1 + s_1^* + \cdots + s_d + s_d^*$, using $\|P\|_{C^*(\mathbf{F}_d)} = 2d$ and $\nu_0(h) = \lim_N \mathbf{E}[\mathrm{tr}_N h(A^N)] = \tau(h(A_F))$ by [Corollary 5.3](#). \square

The remaining ingredient of the proof is to show that $\mathrm{supp} \nu_1 \subseteq [-\|A_F\|, \|A_F\|]$. To this end, it suffices by [Lemma 4.9](#) to understand the exponential growth rate of the moments of ν_1 . We begin by computing a formula for the moments of ν_1 .² Recall that $\mathbf{F}_d^{\mathrm{np}}$ denotes the non-power elements of \mathbf{F}_d (see [Lemma 5.2](#)).

LEMMA 8.2. *For all $p \in \mathbb{N}$, we have*

$$\nu_1(x^p) = -\tau(A_F^p) + \sum_{k=2}^p (\omega(k) - 1) \sum_{v \in \mathbf{F}_d^{\mathrm{np}}} \sum_{i_1, \dots, i_p=1}^{2d} 1_{g_{i_1} \cdots g_{i_p} = v^k}.$$

²In the special setting of random regular graphs, a nonrigorous derivation of an explicit formula for ν_1 (that is, an explicit solution to the Hausdorff moment problem associated to [Lemma 8.2](#)) appears in the physics literature [49]. However, an explicit solution to the moment problem, which is not available in more general situations, is not needed to apply the methods of this paper.

Proof. Note that

$$\begin{aligned} \nu_1(x^p) + \tau(A_{\mathbb{F}}^p) &= \lim_{N \rightarrow \infty} N(\mathbf{E}[\mathrm{tr}_N(A^N)^p] - (1 - \frac{1}{N})\tau(A_{\mathbb{F}}^p)) \\ &= \lim_{N \rightarrow \infty} \sum_{i_1, \dots, i_p=1}^{2d} N(\mathbf{E}[\mathrm{tr}_N S_{i_1}^N \cdots S_{i_p}^N] - (1 - \frac{1}{N})\tau(\lambda(g_{i_1}) \cdots \lambda(g_{i_p}))). \end{aligned}$$

As $\tau(\lambda(g_{i_1}) \cdots \lambda(g_{i_p})) = 1_{g_{i_1} \cdots g_{i_p} = e}$, the terms with $g_{i_1} \cdots g_{i_p} = e$ cancel in the sum as in the proof of [Corollary 5.3](#).

The conclusion now follows from [Lemma 5.2](#) (note that the $k = 1$ term does not appear in the sum as $\omega(1) - 1 = 0$). \square

In the present setting, a simple argument due to Friedman [[26](#), Lemma 2.4] suffices to bound the growth rate of the moments. As a variant of this argument will be needed later in this paper, we recall the proof here.

LEMMA 8.3. *For every $k \geq 2$ and $p \geq 1$, we have*

$$\sum_{v \in \mathbf{F}_d^{\mathrm{np}}} \sum_{i_1, \dots, i_p=1}^{2d} 1_{g_{i_1} \cdots g_{i_p} = v^k} \leq (p+1)^4 \|A_{\mathbb{F}}\|^p.$$

Proof. We would like to argue that if $g_{i_1} \cdots g_{i_p} = v^k$, then there must exist $a < b$ so that $g_{i_1} \cdots g_{i_a} = v$, $g_{i_{a+1}} \cdots g_{i_b} = v$, and $g_{i_{b+1}} \cdots g_{i_p} = v^{k-2}$. But this need not be true: if v is not cyclically reduced, the last letters of v may cancel the first letters of the next repetition of v , and then the cancelled letters need not appear in $g_{i_1} \cdots g_{i_p}$.

To eliminate this ambiguity, we note that any $v \in \mathbf{F}_d^{\mathrm{np}}$ can be uniquely expressed as $v = gwg^{-1}$ with $w \in \mathbf{F}_d^{\mathrm{np}}$, $g \in \mathbf{F}_d$ such that w is cyclically reduced and such that gwg^{-1} is reduced if $g \neq e$. Thus we may write for any $k \geq 2$

$$\begin{aligned} \sum_{v \in \mathbf{F}_d^{\mathrm{np}}} \sum_{i_1, \dots, i_p=1}^{2d} 1_{g_{i_1} \cdots g_{i_p} = v^k} &\leq \sum_{(w,g)} \sum_{0 \leq t_1 \leq t_2 \leq t_3 \leq t_4 \leq p} \sum_{i_1, \dots, i_p=1}^{2d} (1_{g_{i_1} \cdots g_{i_{t_1}} = g} \times \\ &1_{g_{i_{t_1+1}} \cdots g_{i_{t_2}} = w} 1_{g_{i_{t_2+1}} \cdots g_{i_{t_3}} = w} 1_{g_{i_{t_3+1}} \cdots g_{i_{t_4}} = w^{k-2}} 1_{g_{i_{t_4+1}} \cdots g_{i_p} = g^{-1}}), \end{aligned}$$

where the sum is over pairs (w, g) with the above properties. The idea is now to express the indicators as $1_{g_{i_1} \cdots g_{i_t} = v} = \langle \delta_v, \lambda(g_{i_1}) \cdots \lambda(g_{i_t}) \delta_e \rangle$. Substituting this identity into the right-hand side of the above inequality yields

$$\begin{aligned} \sum_{v \in \mathbf{F}_d^{\mathrm{np}}} \sum_{i_1, \dots, i_p=1}^{2d} 1_{g_{i_1} \cdots g_{i_p} = v^k} &\leq \sum_{(w,g)} \sum_{0 \leq t_1 \leq t_2 \leq t_3 \leq t_4 \leq p} (\langle \delta_g, A_{\mathbb{F}}^{t_1} \delta_e \rangle \langle \delta_w, A_{\mathbb{F}}^{t_2-t_1} \delta_e \rangle \times \\ &\langle \delta_w, A_{\mathbb{F}}^{t_3-t_2} \delta_e \rangle \langle \delta_{w^{k-2}}, A_{\mathbb{F}}^{t_4-t_3} \delta_e \rangle \langle \delta_{g^{-1}}, A_{\mathbb{F}}^{p-t_4} \delta_e \rangle) \end{aligned}$$

$$\leq \sum_{0 \leq t_1 \leq t_2 \leq t_3 \leq t_4 \leq p} \|A_F\|^{t_4 - t_3} \left(\sum_{g \in \mathbf{F}_d} \langle \delta_g, A_F^{t_1} \delta_e \rangle \langle \delta_{g^{-1}}, A_F^{p-t_4} \delta_e \rangle \right) \times \\ \left(\sum_{w \in \mathbf{F}_d} \langle \delta_w, A_F^{t_2 - t_1} \delta_e \rangle \langle \delta_w, A_F^{t_3 - t_2} \delta_e \rangle \right),$$

where we used $|\langle \delta_{w^{k-2}}, A_F^{t_4 - t_3} \delta_e \rangle| \leq \|A_F\|^{t_4 - t_3}$. The conclusion now follows readily by applying the Cauchy–Schwarz inequality to the sums over g and w , as

$$\sum_{v \in \mathbf{F}_d} |\langle \delta_v, A_F^t \delta_e \rangle|^2 = \|A_F^t \delta_e\|^2 \leq \|A_F\|^{2t}$$

for any $t \geq 0$. \square

COROLLARY 8.4. $\text{supp } \nu_1 \subseteq [-\|A_F\|, \|A_F\|]$.

Proof. Lemmas 8.2 and 8.3 imply that $|\nu_1(x^p)| \leq (1 + p^2(p+1)^4) \|A_F\|^p$ for all $p \geq 1$, so that the conclusion follows from Lemma 4.9. \square

We can now complete the proof of Theorem 3.4.

Proof of Theorem 3.4. Let χ be the test function provided by Lemma 4.10 with $m = 8$, $K = 2d$, $\rho = \|A_F\| = 2\sqrt{2d-1}$. As $\chi(x) = 0$ for $|x| \leq \|A_F\| + \frac{\varepsilon}{2}$, we have $\tau(\chi(A_F)) = 0$ and $\nu_1(\chi) = 0$ by Corollary 8.4. Thus Lemmas 8.1 and 4.10 yield

$$\mathbf{E}[\text{tr}_N \chi(A^N)] \lesssim \frac{(\log d)^8}{N^2} \beta_* \left(\frac{d}{\varepsilon} \right)^{8+1/\beta_*}$$

for all $\varepsilon < 2d - 2\sqrt{2d-1}$ and $\beta_* \geq 1$.

To conclude, we note that as $\chi(x) \geq 1$ for $|x| \geq \|A_F\| + \varepsilon$, we have

$$\mathbf{P}[\|A^N\| \geq \|A_F\| + \varepsilon] \leq \mathbf{P}[\text{Tr } \chi(A^N) \geq 1] \lesssim \frac{1}{N} \left(\frac{d \log d}{\varepsilon} \right)^8 \log \left(\frac{2ed}{\varepsilon} \right)$$

for $\varepsilon < 2d - 2\sqrt{2d-1}$, where we chose $\beta_* = 1 + \log(\frac{2d}{\varepsilon})$. \square

8.2. *Proof of Theorem 3.6.* The proof is based on results of Puder [61]. Let us begin by synthesizing the parts of [61] that we need here.

LEMMA 8.5. *Define ρ_m as in Theorem 3.6. Then for any $2 \leq m \leq d$, we have*

$$\limsup_{p \rightarrow \infty} |\nu_m(x^p)|^{1/p} \leq \rho_m.$$

Proof. Fix a word $w \in \mathbf{W}_d$ of length q that does not reduce to the identity. By Lemmas 5.1 and 5.2, we can write for all sufficiently large N

$$N \mathbf{E}[\text{tr}_N w(S^N)] = \sum_{s=0}^{\infty} \frac{b_s(w)}{N^s},$$

as a rational function is analytic away from its poles. It is shown in [61, §5.5] that $b_s(w) = 0$ for $s \leq \pi(w) - 2$, that $b_{\pi(w)-1}(w) = |\text{Crit}(w)|$, and that $|b_s(w)| \leq q^{2(s+1)}$ for $s \geq \pi(w)$, where $\pi(w) \in \{0, \dots, d\} \cup \{\infty\}$ denotes the primitivity rank of w . We refer to [61, §2.2] for precise definitions of $\pi(w)$ and $\text{Crit}(w)$, which are not needed in the following argument: the only properties we will use are that $\pi(w) = 0$ if and only if $w = e$, and that $|\text{Crit}(w)| \geq 1$ when $\pi(w) \neq \infty$.

Theorem 6.1 with $m = d + 1$ and $P(\mathbf{s}, \mathbf{s}^*) = s_1 + s_1^* + \dots + s_d + s_d^*$ yields

$$\begin{aligned} \sum_{i_1, \dots, i_p=1}^{2d} N \mathbf{E}[\text{tr}_N S_{i_1}^N \cdots S_{i_p}^N] 1_{g_{i_1} \cdots g_{i_p} \neq e} &= N (\mathbf{E}[\text{tr}_N (A^N)^p] - (1 - \frac{1}{N}) \nu_0(x^p)) \\ &= \nu_0(x^p) + \sum_{i=0}^{d-1} \frac{\nu_{i+1}(x^p)}{N^i} + O\left(\frac{1}{N^d}\right) \end{aligned}$$

for $N \rightarrow \infty$ with p, d fixed (cf. the proof of [Lemma 8.2](#)). It follows that

$$\nu_m(x^p) = \sum_{i_1, \dots, i_p=1}^{2d} b_{m-1}(g_{i_1} \cdots g_{i_p}) 1_{g_{i_1} \cdots g_{i_p} \neq e} = \sum_{r=1}^m \sum_{w \in \mathcal{W}_p^r} b_{m-1}(w)$$

for $m \geq 2$, where $\mathcal{W}_p^r = \{w = g_{i_1} \cdots g_{i_p} : 1 \leq i_1, \dots, i_p \leq 2d, \pi(w) = r\}$. Thus

$$|\nu_m(x^p)| \leq mp^{2m} \max_{r=1, \dots, m} \sum_{w \in \mathcal{W}_p^r} |\text{Crit}(w)|,$$

where we used that $|\text{Crit}(w)| \geq 1$ for $\pi(w) \neq \infty$. The conclusion now follows from the second (unnumbered) theorem in [61, §2.4]. \square

COROLLARY 8.6. *Let $d \geq 2$. Then $\text{supp } \nu_m \subseteq [-\rho_m, \rho_m]$ for all $0 \leq m \leq d$.*

Proof. For $m = 0, 1$ the follows from $\|A_F\| = 2\sqrt{2d-1}$ and [Corollary 8.4](#), while for $m = 2, \dots, d$ this follows from [Lemmas 8.5](#) and [4.9](#). \square

We can now complete the proof of [Theorem 3.6](#).

Proof of Theorem 3.6. Fix $m_* \leq m \leq d - 1$. Let χ be the test function provided by [Lemma 4.10](#) with $m \leftarrow 4(m+1)$, $K \leftarrow 2d$, $\rho \leftarrow \rho_m$. Then $\nu_i(\chi) = 0$ for $i \leq m$ by [Corollary 8.6](#). Thus [Theorem 7.1](#) with $h \leftarrow \chi$, $m \leftarrow m + 1$ and [Lemma 4.10](#) yield

$$\mathbf{P}[\|A^N\| \geq \rho_m + \varepsilon] \leq \mathbf{P}[\text{Tr } \chi(A^N) \geq 1] \lesssim \frac{C_d}{N^m} \frac{1}{\varepsilon^{4(m+1)}} \log\left(\frac{2e}{\varepsilon}\right),$$

where we chose $\beta_* = 1 + \log(\frac{2}{\varepsilon})$ (note that $\beta_* > 1$ as $\varepsilon < \rho_{m+1} - \rho_m \leq 2$).

For the lower bound, it is shown in the proof of [27, Theorem 2.11] that

$$\mathbf{P}[\|A^N\| \geq \alpha'] \geq (1 - o(1)) N^{1-m}$$

for all $m > m_*$ and $\alpha' < \rho_m$. Choosing $\alpha' = \rho_{m-1} + \varepsilon$ completes the proof. \square

9. Strong convergence of random permutation matrices

The aim of this section is to prove [Theorem 3.9](#). Most of the proof is identical to that of [Theorem 3.4](#). The only difficulty in the present setting is to adapt the argument of [Lemma 8.3](#) for bounding $\text{supp } \nu_1$. This is not entirely straightforward, because the proof of [Lemma 8.3](#) relied on an overcounting argument which is not applicable to general polynomials. Nonetheless, we will show that a more careful implementation of the idea behind the proof can avoid this issue.

Throughout the proof, we fix a polynomial P as in [Theorem 3.9](#), and define

$$X^N := P(\mathbf{S}^N, \mathbf{S}^{N*}), \quad X_F := P(\mathbf{s}, \mathbf{s}^*).$$

To simplify the proof, we begin by applying a linearization trick of Pisier [\[59\]](#) in order to factor X_F into polynomials of degree one.

LEMMA 9.1. *For any $\varepsilon > 0$, there exist $q \in \mathbb{N}$ and operators*

$$X_j := \sum_{i=0}^{2d} A_{j,i} \otimes \lambda(g_i), \quad j = 1, \dots, q$$

with matrix coefficients $A_{j,i}$ of dimensions $D_j \times D_{j+1}$ with $D_1 = D_{q+1} = D$, so that $X_F = X_1 \cdots X_q$ and $\|X_j\| \leq (\|X_F\| + \varepsilon)^{1/q}$ for all j .

Proof. This is an immediate consequence of [\[59, Theorem 1\]](#). □

In what follows, we fix $\varepsilon > 0$ and a factorization of X_F as in [Lemma 9.1](#). We will also define X_j^N by replacing $\lambda(g_i) \leftarrow S_i^N$ in the definition of X_j . Note that as g_1, \dots, g_d are free, $X_F = X_1 \cdots X_q$ implies that $X^N = X_1^N \cdots X_q^N$.

It will be convenient to extend the indexing of the above operators cyclically: we will define $X_j := X_{((j-1) \bmod q) + 1}$ for all $j \in \mathbb{N}$. This implies, for example, that $X_F^p = X_1 X_2 \cdots X_{pq}$. We similarly define $A_{j,i} := A_{((j-1) \bmod q) + 1, i}$ for $j \in \mathbb{N}$.

9.1. *The first-order support.* In the present setting, the moments of the linear functionals ν_0 and ν_1 in [Theorem 6.1](#) satisfy

$$\begin{aligned} \nu_0(x^p) &= \lim_{N \rightarrow \infty} \mathbf{E}[\text{tr}_{DN} (X^N)^p] = \tau(X_F^p), \\ \nu_1(x^p) &= \lim_{N \rightarrow \infty} N \left(\mathbf{E}[\text{tr}_{DN} (X^N)^p] - \tau(X_F^p) \right). \end{aligned}$$

We begin by writing an expression for $\nu_1(x^p)$.

LEMMA 9.2. *For every $p \in \mathbb{N}$, we have*

$$\nu_1(x^p) = -\tau(X_F^p) + \sum_{k=2}^{pq} (\omega(k) - 1) \sum_{v \in \mathbf{F}_d^{np}} \sum_{i_1, \dots, i_{pq}=0}^{2d} a_{i_1, \dots, i_{pq}} \mathbf{1}_{g_{i_1} \cdots g_{i_{pq}} = v^k},$$

where we define

$$a_{i_1, \dots, i_{pq}} := \text{tr}_D (A_{1, i_1} A_{2, i_2} \cdots A_{pq, i_{pq}}).$$

Proof. The proof is identical to that of [Lemma 8.2](#). \square

We would like to repeat the proof of [Lemma 8.3](#) in the present setting. Recall that the idea of the proof is to write $v^k = g w w w^{k-2} g^{-1}$ where w is cyclically reduced, and then to reason that any word $g_{i_1} \cdots g_{i_t}$ that reduces to v^k is a concatenation of words $g_{i_1} \cdots g_{i_{t_1}} = g$, $g_{i_{t_1+1}} \cdots g_{i_{t_2}} = w$, $g_{i_{t_2+1}} \cdots g_{i_{t_3}} = w$, etc. We can therefore sum over all such concatenations in the expression for ν_1 .

The problem with this argument is that the above decomposition is not unique: for example, $g_1 g_2 g_2^{-1} g_1 = g_1^2$ can be decomposed in two different ways $(g_1, g_2 g_2^{-1} g_1)$ or $(g_1 g_2 g_2^{-1}, g_1)$. Thus when we sum over all possible ways of generating such concatenations, we are overcounting the number of words that reduce to v^k . Unlike in [Lemma 8.3](#), the coefficients $a_{i_1, \dots, i_{pq}}$ can have both positive and negative signs and thus we cannot upper bound the moments by overcounting.

We presently show how a more careful analysis can avoid overcounting. We begin by introducing the following basic notion.

Definition 9.3. For any $v \in \mathbf{F}_d$ and $0 \leq i_1, \dots, i_k \leq 2d$, we write $g_{i_1} \cdots g_{i_k} \triangleq v$ if $g_{i_1} \cdots g_{i_k} = v$ and $g_{i_l} \cdots g_{i_k} \neq v$ for $1 < l \leq k$.

To interpret this notion, one should think of any word $g_{i_1} \cdots g_{i_k}$ as defining a walk in the Cayley graph of \mathbf{F}_d when read from right to left, starting at the identity. Then $g_{i_1} \cdots g_{i_k} \triangleq v$ states that the walk defined by $g_{i_1} \cdots g_{i_k}$ reaches v for the first time at its endpoint. Just as we can write

$$1_{g_{i_1} \cdots g_{i_k} = v} = \langle \delta_v, \lambda(g_{i_1}) \cdots \lambda(g_{i_k}) \delta_e \rangle,$$

the equivalence notion of [Definition 9.3](#) may also be expressed in terms of matrix elements of the left-regular representation.

LEMMA 9.4. For any $0 \leq i_1, \dots, i_k \leq 2d$ and $v \in \mathbf{F}_d$, we have

$$\begin{aligned} 1_{g_{i_1} \cdots g_{i_k} \triangleq v} &= \langle \delta_v, \lambda(g_{i_1}) \cdots \lambda(g_{i_k}) \delta_e \rangle \\ &\quad - \sum_{s=1}^{k-1} \langle \delta_e, \lambda(g_{i_1}) Q \lambda(g_{i_2}) \cdots Q \lambda(g_{i_s}) \delta_e \rangle \langle \delta_v, \lambda(g_{i_{s+1}}) \cdots \lambda(g_{i_k}) \delta_e \rangle, \end{aligned}$$

where we defined $Q := \mathbf{1} - \delta_e \delta_e^*$ (i.e., Q is the orthogonal projection onto $\{\delta_e\}^\perp$).

Proof. One may simply note that term s in the sum is the indicator of the event that $g_{i_1} \cdots g_{i_k} = v$, $g_{i_{s+1}} \cdots g_{i_k} = v$, and $g_{i_t} \cdots g_{i_k} \neq v$ for $1 < t < s + 1$. \square

We now use this identity to express the kind of sum that appears in [Lemma 9.2](#) exactly (without overcounting) in terms of matrix elements of the operators X_j .

LEMMA 9.5. *Fix $v_1, \dots, v_\ell \neq e \in \mathbf{F}_d$ ($\ell \geq 2$) so that $v_1 \cdots v_\ell$ is reduced. Define*

$$\begin{aligned} X_{(t,s)} &:= X_t \cdots X_s, \\ X_{[t,s]} &:= -X_t(\mathbf{1}_{D_{t+1}} \otimes Q)X_{t+1} \cdots (\mathbf{1}_{D_s} \otimes Q)X_s \end{aligned}$$

for $t \leq s$ and $X_{(t,s)} = X_{[t,s]} := \mathbf{1}$ for $t > s$. Then

$$\begin{aligned} &\sum_{i_1, \dots, i_{pq}=0}^{2d} a_{i_1, \dots, i_{pq}} \mathbf{1}_{g_{i_1} \cdots g_{i_{pq}} = v_1 \cdots v_\ell} = \\ &\sum_{x_2, y_2, \dots, x_\ell, y_\ell, x_{\ell+1}} \sum_{1 < t_2 \leq s_2 < \cdots < t_\ell \leq s_\ell \leq pq} \langle e_{x_{\ell+1}} \otimes \delta_{v_1}, X_{(1, t_2-1)} e_{x_2} \otimes \delta_e \rangle \times \\ &\prod_{r=2}^{\ell} \langle e_{x_r} \otimes \delta_e, X_{[t_r, s_r-1]} e_{y_r} \otimes \delta_e \rangle \langle e_{y_r} \otimes \delta_{v_r}, X_{(s_r, t_{r+1}-1)} e_{x_{r+1}} \otimes \delta_e \rangle, \end{aligned}$$

where we let $t_{\ell+1} := pq + 1$.

Proof. Suppose that $g_{i_1} \cdots g_{i_{pq}} = v_1 \cdots v_\ell$. As we assumed that $v_1 \cdots v_\ell$ is reduced, there are unique indices $1 < t_2 < \cdots < t_\ell \leq pq$ so that

$$g_{i_1} \cdots g_{i_{t_2-1}} = v_1, \quad g_{i_{t_r}} \cdots g_{i_{t_{r+1}-1}} \triangleq v_r \quad \text{for } 2 \leq r \leq \ell.$$

Thus we can write

$$\mathbf{1}_{g_{i_1} \cdots g_{i_{pq}} = v_1 \cdots v_\ell} = \sum_{1 < t_2 < \cdots < t_\ell \leq pq} \mathbf{1}_{g_{i_1} \cdots g_{i_{t_2-1}} = v_1} \prod_{r=2}^{\ell} \mathbf{1}_{g_{i_{t_r}} \cdots g_{i_{t_{r+1}-1}} \triangleq v_r}.$$

The conclusion follows by applying [Lemma 9.4](#) to the indicators in this identity and summing over the indices i_1, \dots, i_{pq} . \square

With this identity in hand, we can proceed exactly as in [Lemma 8.3](#). In the following lemma, recall that we chose an arbitrary $\varepsilon > 0$ in [Lemma 9.1](#).

LEMMA 9.6. *For every $k \geq 2$, we have*

$$\left| \sum_{v \in \mathbf{F}_d^{\text{np}}} \sum_{i_1, \dots, i_{pq}=0}^{2d} a_{i_1, \dots, i_{pq}} \mathbf{1}_{g_{i_1} \cdots g_{i_{pq}} = v^k} \right| \leq 2D_{\max}^9 (pq)^8 (\|X_{\mathbf{F}}\| + \varepsilon)^p,$$

where $D_{\max} = \max_j D_j$.

Proof. Suppose first that $k \geq 3$. Let \mathcal{F} be the set of $v \in \mathbf{F}_d^{\text{np}}$ that are cyclically reduced and let $\mathcal{F}' := \mathbf{F}_d^{\text{np}} \setminus \mathcal{F}$. Then every $v \in \mathcal{F}'$ can be uniquely expressed as $v = gwg^{-1}$ with $w \in \mathcal{F}$, $g \in \mathbf{F}_d$ so that gwg^{-1} is reduced.

Applying [Lemma 9.5](#) with $\ell = 5$ and $v_1 \leftarrow g$, $v_2 \leftarrow w$, $v_3 \leftarrow w$, $v_4 \leftarrow w^{k-2}$, $v_5 \leftarrow g^{-1}$ yields

$$\begin{aligned} & \left| \sum_{v \in \mathcal{F}'} \sum_{i_1, \dots, i_{pq}=0}^{2d} a_{i_1, \dots, i_{pq}} 1_{g_{i_1} \dots g_{i_{pq}} = v^k} \right| \leq \\ & \sum_{1 < t_2 \leq s_2 < \dots < t_5 \leq s_5 \leq pq} \|X_{[t_2, s_2-1]}\| \|X_{[t_3, s_3-1]}\| \|X_{[t_4, s_4-1]}\| \|X_{[t_5, s_5-1]}\| \|X_{(s_4, t_5-1)}\| \times \\ & \sum_{x_2, y_2, \dots, x_5, y_5, x_6} \sum_{w \in \mathbf{F}_d} |\langle e_{y_2} \otimes \delta_w, X_{(s_2, t_3-1)} e_{x_3} \otimes \delta_e \rangle \langle e_{y_3} \otimes \delta_w, X_{(s_3, t_4-1)} e_{x_4} \otimes \delta_e \rangle| \times \\ & \sum_{g \in \mathbf{F}_d} |\langle e_{x_6} \otimes \delta_g, X_{(1, t_2-1)} e_{x_2} \otimes \delta_e \rangle \langle e_{y_5} \otimes \delta_{g^{-1}}, X_{(s_5, pq)} e_{x_6} \otimes \delta_e \rangle|. \end{aligned}$$

Using that $\|X_{(t, s-1)}\| \leq (\|X_{\mathbf{F}}\| + \varepsilon)^{(s-t)/q}$ and $\|X_{[t, s-1]}\| \leq (\|X_{\mathbf{F}}\| + \varepsilon)^{(s-t)/q}$ by [Lemma 9.1](#) and $\|Q\| = 1$, and applying Cauchy–Schwarz as in the proof of [Lemma 8.3](#), we can upper bound the right-hand side by $D_{\max}^9(pq)^8(\|X_{\mathbf{F}}\| + \varepsilon)^p$.

If we sum instead over $v \in \mathcal{F}$ on the left-hand side, we can bound in exactly the same manner by applying [Lemma 9.5](#) with $\ell = 3$ and $v_1 \leftarrow w$, $v_2 \leftarrow w$, $v_3 \leftarrow w^{k-2}$. Thus the proof is complete for the case $k \geq 3$. The proof for $k = 2$ is identical, except that we now omit the w^{k-2} terms. \square

COROLLARY 9.7. *The following holds: $\text{supp } \nu_1 \subseteq [-\|X_{\mathbf{F}}\|, \|X_{\mathbf{F}}\|]$.*

Proof. [Lemmas 9.2](#) and [9.6](#) imply that

$$|\nu_1(x^p)| \leq \|X_{\mathbf{F}}\|^p + 2D_{\max}^9(pq)^{10}(\|X_{\mathbf{F}}\| + \varepsilon)^p.$$

The conclusion follows by first applying [Lemma 4.9](#), and then letting $\varepsilon \rightarrow 0$. \square

9.2. *Proof of [Theorem 3.9](#).* The rest of the proof contains no new ideas.

Proof of [Theorem 3.9](#). Let χ be the test function provided by [Lemma 4.10](#) with $m = 8$, $K = \|P\|_{M_d(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}$, and $\rho = \|X_{\mathbf{F}}\|$. Then $\nu_0(\chi) = \nu_1(\chi) = 0$ by [Corollary 9.7](#). Thus [Theorem 7.1](#) with $h \leftarrow \chi$, $m \leftarrow 2$ and [Lemma 4.10](#) yield

$$\mathbf{P}[\|X^N\| \geq \|X_{\mathbf{F}}\| + \varepsilon] \leq \mathbf{P}[\text{Tr } \chi(X^N) \geq 1] \lesssim \frac{D}{N} \left(\frac{Kq_0 \log d}{\varepsilon} \right)^8 \log \left(\frac{eK}{\varepsilon} \right)$$

for all $\varepsilon < K - \|X_{\mathbf{F}}\|$, where we chose $\beta_* = 1 + \log(\frac{K}{\varepsilon})$. \square

10. Stable representations of the symmetric group

The aim of this section is to prove [Theorem 3.14](#). The main issue here is to extend the basic properties of random permutations matrices of [Section 5](#) to general stable representations. With analogues of these properties in place, the remainder of the proof is nearly the same as that of [Theorem 3.9](#).

Throughout this section, we fix a stable representation as in [Section 3.4.2](#) and adopt the notations introduced there. In addition, we will denote by β the degree of the polynomial $\varphi(x_1, x_2^2, \dots, x_r^r)$ and by $\kappa := \sup_{N \geq N_0} \frac{D_N}{N^\alpha}$.

10.1. *Basic properties.* We will deduce the requisite properties from the much more precise results of [\[32\]](#). (In fact, the only facts that will be needed here are relatively elementary, as is explained in [\[45, Remark 31\]](#) and [\[32, §1.5.1\]](#).)

Let $w \in \mathbf{W}_d$ be a word. Then clearly

$$w(\mathbf{\Pi}^N) = \pi_N(w(\boldsymbol{\sigma})),$$

where $\boldsymbol{\sigma} = (\sigma_1, \dots, \sigma_d)$ are i.i.d. uniformly distributed elements of \mathbf{S}_N . The distribution of $w(\boldsymbol{\sigma})$ defines a probability measure $\mathbf{P}_{w,N}$ on \mathbf{S}_N , called the word measure.

LEMMA 10.1. *Let $w \in \mathbf{W}_d$ be any word of length at most q that does not reduce to the identity, and let $N \geq \max\{\beta q, N_0\}$. Then we have*

$$\mathbf{E}[\mathrm{Tr} w(\mathbf{\Pi}^N)] = \frac{f_w(\frac{1}{N})}{g_q(\frac{1}{N})},$$

where

$$g_q(x) := (1-x)^{d_1} (1-2x)^{d_2} \cdots (1-(\beta q-1)x)^{d_{\beta q-1}}$$

with $d_j := \min(d, \lfloor \frac{\beta q}{j+1} \rfloor)$, and f_w, g_q are polynomials of degree at most

$$\beta q(1 + \log d).$$

Proof. We may assume without loss of generality that w is cyclically reduced with length $1 \leq \ell \leq q$, as $\mathrm{Tr} w(\mathbf{\Pi}^N)$ is invariant under cyclic reduction.

The definition of the stable representation implies that

$$(10.1) \quad \mathbf{E}[\mathrm{Tr} w(\mathbf{\Pi}^N)] = \mathbf{E}_{w,N}[\varphi(\xi_1, \dots, \xi_r)].$$

Let $\xi_1^{\alpha_1} \cdots \xi_r^{\alpha_r}$ be a monomial of φ of degree at least one. Then for N sufficiently large, [\[32, Example 3.6, Definition 6.6, and Proposition 6.8\]](#) imply that

$$(10.2) \quad \mathbf{E}_{w,N}[\xi_1^{\alpha_1} \cdots \xi_r^{\alpha_r}] = \sum_{\Gamma} \frac{\left(\frac{1}{N}\right)^{e_{\Gamma}-v_{\Gamma}} \prod_{i=1}^{v_{\Gamma}-1} \left(1 - \frac{i}{N}\right)}{\prod_{j=1}^d \prod_{i=1}^{e_{\Gamma}^j-1} \left(1 - \frac{i}{N}\right)},$$

where the sum is over a certain collection of quotients Γ of the graph consisting of α_1 disjoint cycles of length ℓ , α_2 disjoint cycles of length 2ℓ , etc., whose edges are colored by $\{1, \dots, d\}$. Here we denote by v_{Γ} the number of vertices of Γ , by e_{Γ}^j the number of edges of Γ colored by j , and by $e_{\Gamma} = e_{\Gamma}^1 + \cdots + e_{\Gamma}^d$.

It is immediate from the above construction that $v_{\Gamma} \leq \ell \sum_{i=1}^r i \alpha_i \leq \beta q$, and analogously $e_{\Gamma} \leq \beta q$. The validity of [\(10.2\)](#) for all $N \geq \beta q \geq \max_j e_{\Gamma}^j$ can be read off from the proof of [\[32, Proposition 6.8\]](#). Moreover, we must

have $e_\Gamma - v_\Gamma \geq 0$ for all Γ that appear in the sum, as $\mathbf{E}_{w,N}[\xi_1^{\alpha_1} \cdots \xi_r^{\alpha_r}] = O(1)$ by [32, Theorem 1.3] and the following discussion. Thus (10.2) is a rational function of $\frac{1}{N}$. The remainder of the proof proceeds exactly as in the proof of Lemma 5.1. \square

We now describe the lowest order asymptotics.

LEMMA 10.2. *Fix a word $w \in \mathbf{W}_d$ that does not reduce to the identity, and express its reduction as $w(g_1, \dots, g_d) = v^k$ with $v \in \mathbf{F}_d^{\text{np}}$ and $k \geq 1$. Then*

$$\lim_{N \rightarrow \infty} \mathbf{E}[\text{Tr } w(\mathbf{\Pi}^N)] = \varpi(k)$$

with $\varpi(1) = 0$ and $|\varpi(k)| \leq ck^\beta$ for $k \geq 2$. Here c is a constant that depends on φ .

Proof. Recall that φ is a sum of character polynomials of irreducible representations, and that we assumed that π_N does not contain the trivial representation. The case $k = 1$ therefore follows from Lemma 5.2 for the standard representation and from [32, Corollary 1.7] for all other irreducible components of φ .

For $k \geq 2$, let $\xi_1^{\alpha_1} \cdots \xi_r^{\alpha_r}$ be a monomial of φ . Applying [32, Theorem 1.3] and the following discussion as well as [32, Remark 7.3] yields

$$\lim_{N \rightarrow \infty} \mathbf{E}_{w,N}[\xi_1^{\alpha_1} \cdots \xi_r^{\alpha_r}] = \lim_{N \rightarrow \infty} \mathbf{E}_{v,N}[\xi_k^{\alpha_1} \cdots \xi_{rk}^{\alpha_r}] = \sum_{\substack{\mathcal{P} \in \\ \text{Partitions}(S)}} \prod_{A \in \mathcal{P}} \sum_{j | \gcd(A)} j^{|A|-1},$$

where S is the multiset that contains jk with multiplicity α_j for $j = 1, \dots, r$. By crudely estimating $j \leq \gcd(A) \leq rk$, the right-hand side is bounded by $B_{|S|}(rk)^{|S|}$ where $B_n \leq n!$ denotes the number of partitions of a set with n elements. The conclusion follows as $|S| = \sum_{j=1}^r \alpha_j \leq \beta$ and using (10.1). \square

Note that an immediate consequence of Lemma 10.2 is that the weak convergence as in Corollary 5.3 remains valid in the present setting.

10.2. *Proof of Theorem 3.14.* Fix a self-adjoint noncommutative polynomial $P \in \mathbf{M}_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ of degree q_0 , and let $h \in \mathcal{P}_q$. Then $(\text{Tr} \otimes \text{id})(h(P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}))$ is a linear combination of words $w(\mathbf{\Pi}^N)$ of length at most qq_0 . Summing separately over words that do and do not reduce to the identity yields

$$\mathbf{E}[\text{Tr } h(P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}))] = \frac{\tilde{f}_h(\frac{1}{N})}{g_{qq_0}(\frac{1}{N})} + D_N(\text{Tr} \otimes \tau)(h(P(\mathbf{s}, \mathbf{s}^*)))$$

by Lemma 10.1, where \tilde{f}_h is a polynomial of degree at most $\beta qq_0(1 + \log d)$. Thus

$$\mathbf{E}[\text{tr}_{N^\alpha} h(P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}))] = \Psi_h(\frac{1}{N}) = \frac{f_h(\frac{1}{N})}{g_{qq_0}(\frac{1}{N})},$$

where f_h is the polynomial of degree at most $\beta q q_0(1 + \log d) + \alpha \leq 2\beta q q_0(1 + \log d)$. Here we used that $\frac{D_N}{N^\alpha}$ is a polynomial of $\frac{1}{N}$ of degree α .

We can now repeat the proofs in [sections 6–7](#) nearly verbatim with the replacements $q \leftarrow 2\beta N_0^{1/2} q$ (here we multiply by $N_0^{1/2}$ to ensure $M \geq N_0$ in [Lemma 6.4](#)) and $|\Psi_h(\frac{1}{N})| \leq \frac{DD_N}{N^\alpha} \|h\|_{C^0[-K,K]} \leq \kappa D \|h\|_{C^0[-K,K]}$ to conclude the following:

LEMMA 10.3. *Let $d \geq 2$. Fix P as in [Theorem 3.14](#) and*

$$K = \|P\|_{M_D(\mathbb{C}) \otimes C^*(\mathbf{F}_d)}.$$

Then there exist compactly supported distributions ν_i such that

$$\begin{aligned} & \left| \mathbf{E} \left[\text{tr}_{N^\alpha} h(P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N^*})) \right] - \sum_{i=0}^{\alpha} \frac{\nu_i(h)}{N^i} \right| \\ & \leq \frac{CD (q_0 \log d)^{4(\alpha+1)}}{N^{\alpha+1}} \beta_* \|f^{(4(\alpha+1)+1)}\|_{L^\beta[0,2\pi]} \end{aligned}$$

for all $N \geq N_0$, $\beta > 1$, and $h \in C^\infty(\mathbb{R})$. Here $f(\theta) := h(K \cos \theta)$, $1/\beta_ := 1 - 1/\beta$, and C is a constant that depends on the choice of stable representation.*

Next, we must characterize the supports of the distributions ν_i .

LEMMA 10.4. *The following holds: $\text{supp } \nu_i \subseteq [-\|P(\mathbf{s}, \mathbf{s}^*)\|, \|P(\mathbf{s}, \mathbf{s}^*)\|]$ for $i = 0, \dots, \alpha$.*

Proof. Suppose first that $0 \leq i < \alpha$. As [Lemma 10.2](#) implies that

$$\mathbf{E} \left[\text{tr}_{N^\alpha} h(P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N^*})) \right] = \frac{D_N}{N^\alpha} (\text{Tr} \otimes \tau) \left(h(P(\mathbf{s}, \mathbf{s}^*)) \right) + O\left(\frac{1}{N^\alpha}\right)$$

as $N \rightarrow \infty$ for every $h \in \mathcal{P}$, it follows that there is a constant c_i for every $i < \alpha$ so that $\nu_i(h) = c_i (\text{Tr} \otimes \tau)(h(P(\mathbf{s}, \mathbf{s}^*)))$. The claim follows directly.

We now consider $i = \alpha$. Then we can repeat the proof of [Lemma 9.2](#), but using [Lemma 10.2](#) instead of [Lemma 5.2](#), to obtain the representation

$$\nu_\alpha(x^p) = D_0 \tau(P(\mathbf{s}, \mathbf{s}^*)^p) + \sum_{k=2}^{pq} \varpi(k) \sum_{v \in \mathbf{F}_d^{\text{np}}} \sum_{i_1, \dots, i_{pq}=0}^{2d} a_{i_1, \dots, i_{pq}} \mathbf{1}_{g_{i_1} \dots g_{i_{pq}} = v^k},$$

where $D_0 := \varphi(0, \dots, 0)$ is the constant term in the polynomial $N \mapsto D_N$. We therefore obtain $|\nu_\alpha(x^p)| \lesssim D_{\max}^9 (pq)^{9+\beta} (\|P(\mathbf{s}, \mathbf{s}^*)\| + \varepsilon)^p$ by [Lemmas 9.6](#) and [10.2](#), and the conclusion follows by [Lemma 4.9](#) and as $\varepsilon > 0$ is arbitrary. \square

The rest of the proof follows in the usual manner.

Proof of [Theorem 3.14](#). Let χ be the test function provided by [Lemma 4.10](#) with $m \leftarrow 4(\alpha + 1)$ and $\rho \leftarrow \|P(\mathbf{s}, \mathbf{s}^*)\|$. Then $\nu_i(\chi) = 0$ for all $i \leq \alpha$ by

Lemma 10.4. Thus [Lemma 10.3](#) with $h \leftarrow \chi$, and [Lemma 4.10](#) yield

$$\mathbf{P} \left[\|P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})\| \geq \|P(\mathbf{s}, \mathbf{s}^*)\| + \varepsilon \right] \leq \frac{CD}{N} \left(\frac{Kq_0 \log d}{\varepsilon} \right)^{4(\alpha+1)} \log \left(\frac{eK}{\varepsilon} \right),$$

where we chose $\beta_* = 1 + \log(\frac{K}{\varepsilon})$. \square

Appendix A. Lower bounds

There is a general principle that any random matrix model that satisfies

$$\|P(\mathbf{X}^N, \mathbf{X}^{N*})\| \leq \|P(\mathbf{x}, \mathbf{x}^*)\| + o(1) \quad \text{with probability } 1 - o(1)$$

for every noncommutative polynomial P , automatically also satisfies

$$\|P(\mathbf{X}^N, \mathbf{X}^{N*})\| \geq \|P(\mathbf{x}, \mathbf{x}^*)\| - o(1) \quad \text{with probability } 1 - o(1),$$

provided that the C^* -algebra generated by \mathbf{x} has a unique faithful trace; this can be shown, for example, along the lines of [46, pp.16–19]. Since the latter property holds for the limiting model defined in [Section 3.1](#) (cf. [60]), this general principle applies to all the models considered in this paper.

For this reason, we have focused exclusively on strong convergence upper bounds in the main part of this paper. On the other hand, such a general principle is usually not needed in practice, as in most cases the lower bound already follows easily from *weak* convergence alone. For completeness, we presently provide an elementary proof of the lower bound in the setting of this paper along these lines.

THEOREM A.1. *Let $P \in M_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ be any noncommutative polynomial, and fix any stable representation as in [Section 3.4.2](#). Then*

$$\|P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})\| \geq \|P(\mathbf{s}, \mathbf{s}^*)\| - o(1) \quad \text{with probability } 1 - o(1)$$

as $N \rightarrow \infty$.

Note that all the models in this paper may be viewed as special cases of stable representations, so that [Theorem A.1](#) captures random permutation matrices and random regular graphs as well. The proof of [Theorem A.1](#) could also form the basis for quantitative lower bounds, but we do not pursue this here.

We begin by recalling the basic fact that [Theorem A.1](#) will follow directly once we establish weak convergence in the following form.

LEMMA A.2 (Weak convergence in probability). *In the setting of [Theorem A.1](#),*

$$\mathrm{tr}_{DD_N}(P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})) = (\mathrm{tr}_D \otimes \tau)(P(\mathbf{s}, \mathbf{s}^*)) + o(1)$$

with probability $1 - o(1)$ as $N \rightarrow \infty$ for every self-adjoint $P \in M_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^ \rangle$.*

Let us first use this result to complete the proof of [Theorem A.1](#).

Proof of Theorem A.1. Fix the polynomial P . Then

$$\|P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})\|^{2p} \geq \text{tr}_{DD_N}(|P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})|^{2p}) \geq (\text{tr}_D \otimes \tau)(|P(\mathbf{s}, \mathbf{s}^*)|^{2p}) - o(1)$$

with probability $1 - o(1)$ as $N \rightarrow \infty$ for every $p \in \mathbb{N}$ by Lemma A.2. Moreover,

$$(\text{tr}_D \otimes \tau)(|P(\mathbf{s}, \mathbf{s}^*)|^{2p})^{1/2p} = \|P(\mathbf{s}, \mathbf{s}^*)\| - o(1)$$

as $p \rightarrow \infty$ due to the fact that τ defines a faithful state on the C^* -algebra $C_{\text{red}}^*(\mathbf{F}_d)$ that is generated by \mathbf{s} [54, Prop. 3.12]. The conclusion follows. \square

The only subtlety in implementing this idea is that the form of weak convergence that follows from Lemma 10.2 only yields convergence in expectation

$$(A.1) \quad \mathbf{E}[\text{tr}_{DD_N} P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})] = (\text{tr}_D \otimes \tau)(P(\mathbf{s}, \mathbf{s}^*)) + o(1),$$

as opposed to convergence in probability as in Lemma A.2. We must therefore still show concentration around the mean. The latter can be deduced from (A.1) using a minor additional argument. Note that (A.1) does not require P to be self-adjoint.

Proof of Lemma A.2. Since every stable representation is the direct sum of irreducible stable representations (cf. Section 3.4.1), we can assume in the rest of the proof that π_N is irreducible. Moreover, given any $P \in M_D(\mathbb{C}) \otimes \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$,

$$\text{tr}_{DD_N} P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}) = \frac{1}{D} \sum_{i=1}^D \text{tr}_{D_N} P_{ii}(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}),$$

where $P_{ii} \in \mathbb{C}\langle \mathbf{s}, \mathbf{s}^* \rangle$ are the diagonal elements of P with respect to the matrix coefficients. We can therefore assume in the rest of the proof that $D = 1$.

Let $\Pi_0^N = \pi_N(\sigma_0)$, where σ_0 is a uniformly distributed random element of \mathbf{S}_N that is independent of $\sigma_1, \dots, \sigma_d$. Moreover, let $s_0 = \lambda(g_0)$ where g_0, \dots, g_d are free generators of \mathbf{F}_{d+1} . It follows from Schur's lemma that

$$\mathbf{E}[\Pi_0^N M \Pi_0^{N*}] = \text{tr}_{D_N}(M) \mathbf{1}$$

for every $M \in M_{D_N}(\mathbb{C})$. Therefore

$$\mathbf{E}[\text{tr}_{D_N} \Pi_0^N P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}) \Pi_0^{N*} P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})] = \mathbf{E}[(\text{tr}_{D_N} P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*}))^2].$$

Moreover, it is readily verified using the definitions in Section 3.1 that

$$\tau(s_0 P(\mathbf{s}, \mathbf{s}^*) s_0^* P(\mathbf{s}, \mathbf{s}^*)) = (\tau(P(\mathbf{s}, \mathbf{s}^*)))^2.$$

Since the objects inside the traces on the left-hand side are noncommutative polynomials in $d + 1$ variables and as P is self-adjoint, applying (A.1) yields

$$\lim_{N \rightarrow \infty} \text{Var}(\text{tr}_{D_N} P(\mathbf{\Pi}^N, \mathbf{\Pi}^{N*})) = 0.$$

Given that convergence in expectation follows from (A.1) and that the variance converges to zero, convergence in probability follows immediately. \square

Acknowledgments. We are grateful to Charles Bordenave, Benoît Collins, Michael Magee, Doron Puder, and Mikael de la Salle for very helpful discussions, and to Nick Cook, Srivatsav Kunnawalkam Elayavalli, Joel Friedman, Sidhanth Mohanty, Peter Sarnak, Nikhil Srivastava, Pierre Youssef, and Yizhe Zhu for helpful comments. We are especially indebted to Michael Magee for suggesting the application to stable representations, and for patiently explaining what they are. RvH thanks Peter Sarnak for organizing a stimulating seminar on strong convergence and related topics at Princeton in Fall 2023. Finally, we are grateful to the referee for suggestions that helped us improve the presentation of the paper.

CFC was supported by a Simons-CIQC postdoctoral fellowship through NSF grant QLCI-2016245. JGV was supported by NSF grant FRG-1952777, Caltech IST, and a Baer–Weiss postdoctoral fellowship. JAT was supported by the Caltech Carver Mead New Adventures Fund, NSF grant FRG-1952777, and ONR grants BRC-N00014-18-1-2363 and N00014-24-1-2223. RvH was supported by NSF grants DMS-2054565 and DMS-2347954.

References

- [1] S. AARONSON, The polynomial method in quantum and classical computing, in *Proc. 49th IEEE Symposium on Foundations of Computer Science*, IEEE, 2008, p. 3. <https://doi.org/10.1109/FOCS.2008.91>.
- [2] A. S. BANDEIRA, M. T. BOEDIHARDJO, and R. VAN HANDEL, Matrix concentration inequalities and free probability, *Invent. Math.* **234** no. 1 (2023), 419–487. [MR 4635836. Zbl 1527.60020. https://doi.org/10.1007/s00222-023-01204-6](https://doi.org/10.1007/s00222-023-01204-6).
- [3] R. BAUERSCHMIDT, J. HUANG, A. KNOWLES, and H.-T. YAU, Edge rigidity and universality of random regular graphs of intermediate degree, *Geom. Funct. Anal.* **30** no. 3 (2020), 693–769. [MR 4135670. Zbl 1453.05117. https://doi.org/10.1007/s00039-020-00538-0](https://doi.org/10.1007/s00039-020-00538-0).
- [4] S. BELINSCHI and M. CAPITAINE, Strong convergence of tensor products of independent G.U.E. matrices, 2024. [arXiv 2205.07695](https://arxiv.org/abs/2205.07695).
- [5] C. BORDENAVE, A new proof of Friedman’s second eigenvalue theorem and its extension to random lifts, *Ann. Sci. Éc. Norm. Supér. (4)* **53** no. 6 (2020), 1393–1439. [MR 4203039. Zbl 1462.05324. https://doi.org/10.24033/asens.2450](https://doi.org/10.24033/asens.2450).
- [6] C. BORDENAVE, D. CHAFAÏ, and D. GARCÍA-ZELADA, Convergence of the spectral radius of a random matrix through its characteristic polynomial, *Probab. Theory Related Fields* **182** no. 3-4 (2022), 1163–1181. [MR 4408512. Zbl 1492.30022. https://doi.org/10.1007/s00440-021-01079-9](https://doi.org/10.1007/s00440-021-01079-9).
- [7] C. BORDENAVE and B. COLLINS, Eigenvalues of random lifts and polynomials of random permutation matrices, *Ann. of Math. (2)* **190** no. 3 (2019), 811–875. [MR 4024563. Zbl 1446.60004. https://doi.org/10.4007/annals.2019.190.3.3](https://doi.org/10.4007/annals.2019.190.3.3).
- [8] C. BORDENAVE and B. COLLINS, Norm of matrix-valued polynomials in random unitaries and permutations, 2024. [arXiv 2304.05714](https://arxiv.org/abs/2304.05714).

- [9] C. BORDENAVE and B. COLLINS, Strong asymptotic freeness for independent uniform variables on compact groups associated to nontrivial representations, *Invent. Math.* **237** no. 1 (2024), 221–273. MR 4756991. Zbl 1553.46064. <https://doi.org/10.1007/s00222-024-01259-z>.
- [10] P. BORWEIN and T. ERDÉLYI, *Polynomials and Polynomial Inequalities, Graduate Texts in Mathematics* **161**, Springer-Verlag, New York, 1995. MR 1367960. Zbl 0840.26002. <https://doi.org/10.1007/978-1-4612-0793-1>.
- [11] J. BOURGAIN and L. TZAFRIRI, On a problem of Kadison and Singer, *J. Reine Angew. Math.* **420** (1991), 1–43. MR 1124564. Zbl 0729.47028. <https://doi.org/10.1515/crll.1991.420.1>.
- [12] T. BRAILOVSKAYA and R. VAN HANDEL, Universality and sharp matrix concentration inequalities, *Geom. Funct. Anal.* **34** no. 6 (2024), 1734–1838. MR 4823211. Zbl 1555.60017. <https://doi.org/10.1007/s00039-024-00692-9>.
- [13] E. CASSIDY, Random permutations acting on k -tuples have near-optimal spectral gap for $k = \text{poly}(n)$, 2025. arXiv 2412.13941.
- [14] C.-F. CHEN, J. DOCTER, M. XU, A. BOULAND, F. G. S. L. BRANDÃO, and P. HAYDEN, Efficient unitary designs from random sums and permutations, in 2024 *IEEE 65th Annual Symposium on Foundations of Computer Science—FOCS 2024*, IEEE Computer Soc., Los Alamitos, CA, 2024, pp. 476–484. MR 4849233.
- [15] C.-F. CHEN, J. DOCTER, M. XU, A. BOULAND, F. G. S. L. BRANDÃO, and P. HAYDEN, Efficient unitary designs from random sums and permutations, in 2024 *IEEE 65th Annual Symposium on Foundations of Computer Science—FOCS 2024*, IEEE Computer Soc., Los Alamitos, CA, 2024, pp. 476–484. MR 4849233.
- [16] E. W. CHENEY, *Introduction to Approximation Theory*, AMS Chelsea Publishing, Providence, RI, 1998, Reprint of the second (1982) edition. MR 1656150. Zbl 0912.41001.
- [17] M. D. CHOI, The full C^* -algebra of the free group on two generators, *Pacific J. Math.* **87** no. 1 (1980), 41–48. MR 0590864. Zbl 0463.46047. <https://doi.org/10.2140/pjm.1980.87.41>.
- [18] T. CHURCH, J. S. ELLENBERG, and B. FARB, FI-modules and stability for representations of symmetric groups, *Duke Math. J.* **164** no. 9 (2015), 1833–1910. MR 3357185. Zbl 1339.55004. <https://doi.org/10.1215/00127094-3120274>.
- [19] B. COLLINS, Moment methods on compact groups: Weingarten calculus and its applications, in *ICM—International Congress of Mathematicians. Vol. 4. Sections 5–8*, EMS Press, Berlin, 2023, pp. 3142–3164. MR 4680355. Zbl 1544.43001.
- [20] B. COLLINS, A. GUIONNET, and F. PARRAUD, On the operator norm of non-commutative polynomials in deterministic matrices and iid GUE matrices, *Camb. J. Math.* **10** no. 1 (2022), 195–260. MR 4445344. Zbl 1487.46073. <https://doi.org/10.4310/cjm.2022.v10.n1.a3>.
- [21] B. COLLINS, S. MATSUMOTO, and J. NOVAK, The Weingarten calculus, *Notices Amer. Math. Soc.* **69** no. 5 (2022), 734–745. MR 4415894. Zbl 1527.60007. <https://doi.org/10.1090/noti2474>.

- [22] D. COPPERSMITH and T. J. RIVLIN, The growth of polynomials bounded at equally spaced points, *SIAM J. Math. Anal.* **23** no. 4 (1992), 970–983. MR 1166569. Zbl 0769.26003. <https://doi.org/10.1137/0523054>.
- [23] S. COSTE, G. LAMBERT, and Y. ZHU, The characteristic polynomial of sums of random permutations and regular digraphs, *Int. Math. Res. Not. IMRN* no. 3 (2024), 2461–2510. MR 4702282. Zbl 1554.60012. <https://doi.org/10.1093/imrn/rnad182>.
- [24] R. ESTRADA and R. P. KANWAL, Moment sequences for a certain class of distributions, *Complex Variables Theory Appl.* **9** no. 1 (1987), 31–39. MR 0916914. Zbl 0643.46032. <https://doi.org/10.1080/17476938708814247>.
- [25] B. FARB, Representation stability, in *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II*, Kyung Moon Sa, Seoul, 2014, pp. 1173–1196. MR 3728657. Zbl 1373.55006.
- [26] J. FRIEDMAN, Relative expanders or weakly relatively Ramanujan graphs, *Duke Math. J.* **118** no. 1 (2003), 19–35. MR 1978881. Zbl 1035.05058. <https://doi.org/10.1215/S0012-7094-03-11812-8>.
- [27] J. FRIEDMAN, A proof of Alon’s second eigenvalue conjecture and related problems, *Mem. Amer. Math. Soc.* **195** no. 910 (2008), viii+100. MR 2437174. Zbl 1177.05070. <https://doi.org/10.1090/memo/0910>.
- [28] J. FRIEDMAN, A. JOUX, Y. ROICHMAN, J. STERN, and J.-P. TILLICH, The action of a few random permutations on r -tuples and an application to cryptography, in *STACS 96 (Grenoble, 1996)*, *Lecture Notes in Comput. Sci.* **1046**, Springer, Berlin, 1996, pp. 375–386. MR 1462111. Zbl 1380.94089. https://doi.org/10.1007/3-540-60922-9_31.
- [29] J. FRIEDMAN and D.-E. KOHLER, The relativized second eigenvalue conjecture of Alon, 2014. [arXiv 1403.3462](https://arxiv.org/abs/1403.3462).
- [30] U. HAAGERUP, H. SCHULTZ, and S. THORBJØRNSEN, A random matrix approach to the lack of projections in $C_{\text{red}}^*(\mathbb{F}_2)$, *Adv. Math.* **204** no. 1 (2006), 1–83. MR 2233126. Zbl 1109.15020. <https://doi.org/10.1016/j.aim.2005.05.008>.
- [31] U. HAAGERUP and S. THORBJØRNSEN, A new application of random matrices: $\text{Ext}(C_{\text{red}}^*(\mathbb{F}_2))$ is not a group, *Ann. of Math. (2)* **162** no. 2 (2005), 711–775. MR 2183281. Zbl 1103.46032. <https://doi.org/10.4007/annals.2005.162.711>.
- [32] L. HANANY and D. PUDER, Word measures on symmetric groups, *Int. Math. Res. Not. IMRN* no. 11 (2023), 9221–9297. MR 4597208. Zbl 1522.20016. <https://doi.org/10.1093/imrn/rnac084>.
- [33] M. B. HASTINGS and A. W. HARROW, Classical and quantum tensor product expanders, *Quantum Inf. Comput.* **9** no. 3-4 (2009), 336–360. MR 2553116. Zbl 1171.81329.
- [34] B. HAYES, A random matrix approach to the Peterson-Thom conjecture, *Indiana Univ. Math. J.* **71** no. 3 (2022), 1243–1297. MR 4448584. Zbl 1503.46049. <https://doi.org/10.1512/iumj.2022.71.9386>.
- [35] B. HAYES, D. JEKEL, and S. KUNNAWALKAM ELAYAVALLI, Consequences of the random matrix solution to the Peterson-Thom conjecture, *Anal. PDE* **18** no. 7

- (2025), 1805–1834. MR 4930595. Zbl 08061682. <https://doi.org/10.2140/apde.2025.18.1805>.
- [36] Y. HE, Spectral gap and edge universality of dense random regular graphs, *Comm. Math. Phys.* **405** no. 8 (2024), Paper No. 181, 40. MR 4777082. Zbl 1545.05200. <https://doi.org/10.1007/s00220-024-05063-x>.
- [37] W. HIDE, Effective lower bounds for spectra of random covers and random unitary bundles, *Israel J. Math.* (2025), To appear. arXiv 2305.0458.
- [38] W. HIDE and M. MAGEE, Near optimal spectral gaps for hyperbolic surfaces, *Ann. of Math.* (2) **198** no. 2 (2023), 791–824. MR 4635304. Zbl 1553.58014. <https://doi.org/10.4007/annals.2023.198.2.6>.
- [39] L. HÖRMANDER, *The Analysis of Linear Partial Differential Operators. I, Classics in Mathematics*, Springer-Verlag, Berlin, 2003, Distribution theory and Fourier analysis, Reprint of the second (1990) edition [Springer, Berlin; MR1065993 (91m:35001a)]. MR 1996773. Zbl 1028.35001. <https://doi.org/10.1007/978-3-642-61497-2>.
- [40] J. HUANG, T. MCKENZIE, and H.-T. YAU, Optimal eigenvalue rigidity of random regular graphs, 2024. arXiv 2405.12161.
- [41] J. HUANG, T. MCKENZIE, and H.-T. YAU, Ramanujan property and edge universality of random regular graphs, 2025. arXiv 2412.20263.
- [42] J. HUANG and H.-T. YAU, Spectrum of random d -regular graphs up to the edge, *Comm. Pure Appl. Math.* **77** no. 3 (2024), 1635–1723. MR 4692880. Zbl 1531.05220. <https://doi.org/10.1002/cpa.22176>.
- [43] H. KESTEN, Symmetric random walks on groups, *Trans. Amer. Math. Soc.* **92** (1959), 336–354. MR 0109367. Zbl 0092.33503. <https://doi.org/10.2307/1993160>.
- [44] F. LEHNER, Computing norms of free operators with matrix coefficients, *Amer. J. Math.* **121** no. 3 (1999), 453–486. MR 1738412. Zbl 0929.22004. <https://doi.org/10.1353/ajm.1999.0022>.
- [45] N. LINIAL and D. PUDER, Word maps and spectra of random graph lifts, *Random Structures Algorithms* **37** no. 1 (2010), 100–135. MR 2674623. Zbl 1242.60011. <https://doi.org/10.1002/rsa.20304>.
- [46] L. LOUDER and M. MAGEE, Strongly convergent unitary representations of limit groups, *J. Funct. Anal.* **288** no. 6 (2025), Paper No. 110803, 28. MR 4847195. Zbl 1560.22015. <https://doi.org/10.1016/j.jfa.2024.110803>.
- [47] R. C. LYNDON and P. E. SCHUPP, *Combinatorial Group Theory*, 1977 ed., *Classics in Mathematics*, Springer-Verlag, Berlin, 2001, Reprint of the 1977 edition. MR 1812024. Zbl 0997.20037. <https://doi.org/10.1007/978-3-642-61896-3>.
- [48] M. MAGEE and J. THOMAS, Strongly convergent unitary representations of right-angled artin groups, 2023. arXiv 2308.00863.
- [49] F. L. METZ, G. PARISI, and L. LEUZZI, Finite-size corrections to the spectrum of regular random graphs: An analytical solution, *Phys. Rev. E* **90** no. 5 (2014), 11. Zbl arXiv:1403.2582. <https://doi.org/10.1103/PhysRevE.90.052109>.
- [50] J. A. MINGO and R. SPEICHER, *Free Probability and Random Matrices, Fields Institute Monographs* **35**, Springer, New York; Fields Institute for Research in

- Mathematical Sciences, Toronto, ON, 2017. [MR 3585560](#). [Zbl 1387.60005](#). <https://doi.org/10.1007/978-1-4939-6942-5>.
- [51] S. MOHANTY, R. O'DONNELL, and P. PAREDES, Explicit near-Ramanujan graphs of every degree, in *STOC '20—Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, ACM, New York, 2020, pp. 510–523. [MR 4141778](#). [Zbl 1509.68206](#).
- [52] A. NICA, Asymptotically free families of random unitaries in symmetric groups, *Pacific J. Math.* **157** no. 2 (1993), 295–310. [MR 1197059](#). [Zbl 0739.46051](#). <https://doi.org/10.2140/pjm.1993.157.295>.
- [53] A. NICA, On the number of cycles of given length of a free word in several random permutations, *Random Structures Algorithms* **5** no. 5 (1994), 703–730. [MR 1300595](#). [Zbl 0808.60018](#). <https://doi.org/10.1002/rsa.3240050506>.
- [54] A. NICA and R. SPEICHER, *Lectures on the Combinatorics of Free Probability*, *London Mathematical Society Lecture Note Series* **335**, Cambridge University Press, Cambridge, 2006. [MR 2266879](#). [Zbl 1133.60003](#). <https://doi.org/10.1017/CBO9780511735127>.
- [55] A. NILI, On the second eigenvalue of a graph, *Discrete Math.* **91** no. 2 (1991), 207–210. [MR 1124768](#). [Zbl 0771.05064](#). [https://doi.org/10.1016/0012-365X\(91\)90112-F](https://doi.org/10.1016/0012-365X(91)90112-F).
- [56] R. O'DONNELL and X. WU, Explicit near-fully X-Ramanujan graphs, in *2020 IEEE 61st Annual Symposium on Foundations of Computer Science*, IEEE Computer Soc., Los Alamitos, CA, 2020, pp. 1045–1056. [MR 4232109](#). [Zbl 08079033](#).
- [57] F. PARRAUD, On the operator norm of non-commutative polynomials in deterministic matrices and iid Haar unitary matrices, *Probab. Theory Related Fields* **182** no. 3-4 (2022), 751–806. [MR 4408503](#). [Zbl 1487.60011](#). <https://doi.org/10.1007/s00440-021-01101-0>.
- [58] F. PARRAUD, Asymptotic expansion of smooth functions in polynomials in deterministic matrices and iid GUE matrices, *Comm. Math. Phys.* **399** no. 1 (2023), 249–294. [MR 4567374](#). [Zbl 1520.46035](#). <https://doi.org/10.1007/s00220-022-04551-2>.
- [59] G. PISIER, On a linearization trick, *Enseign. Math.* **64** no. 3-4 (2018), 315–326. [MR 3987147](#). [Zbl 1462.46063](#). <https://doi.org/10.4171/LEM/64-3/4-5>.
- [60] R. T. POWERS, Simplicity of the C^* -algebra associated with the free group on two generators, *Duke Math. J.* **42** no. 2 (1975), 151–156. [MR 0374334](#). [Zbl 0342.46046](#). <https://doi.org/10.1215/S0012-7094-75-04213-1>.
- [61] D. PUDER, Expansion of random graphs: new proofs, new results, *Invent. Math.* **201** no. 3 (2015), 845–908. [MR 3385636](#). [Zbl 1320.05115](#). <https://doi.org/10.1007/s00222-014-0560-x>.
- [62] I. RIVIN and N. T. SARDARI, Quantum chaos on random Cayley graphs of $SL_2[\mathbb{Z}/p\mathbb{Z}]$, *Exp. Math.* **28** no. 3 (2019), 328–341. [MR 3985838](#). [Zbl 1420.81013](#). <https://doi.org/10.1080/10586458.2017.1403982>.
- [63] A. SARID, The spectral gap of random regular graphs, *Random Structures Algorithms* **63** no. 2 (2023), 557–587. [MR 4622336](#). [Zbl 1525.05174](#). <https://doi.org/10.1002/rsa.21150>.

- [64] H. SCHULTZ, Non-commutative polynomials of independent Gaussian random matrices. The real and symplectic cases, *Probab. Theory Related Fields* **131** no. 2 (2005), 261–309. MR 2117954. Zbl 1085.46045. <https://doi.org/10.1007/s00440-004-0366-7>.
- [65] A. SONG, Random harmonic maps into spheres, 2025. [arXiv 2402.10287](https://arxiv.org/abs/2402.10287).
- [66] V. VU, Random discrete matrices, in *Horizons of Combinatorics*, *Bolyai Soc. Math. Stud.* **17**, Springer, Berlin, 2008, pp. 257–280. MR 2432537. Zbl 1154.15024. https://doi.org/10.1007/978-3-540-77200-2_13.
- [67] A. ZYGMUND, *Trigonometric Series. Vol. I, II*, Third ed., *Cambridge Mathematical Library*, Cambridge University Press, Cambridge, 2002, With a foreword by Robert A. Fefferman. MR 1963498. Zbl 1084.42003.

(Received: June 26, 2024)

(Revised: February 11, 2025)

CHI-FANG CHEN

EECS, University of California, Berkeley, CA, US and Center for Theoretical Physics,
Massachusetts Institute of Technology, Cambridge, MA, USA

E-mail: achifchen@gmail.com

<https://orcid.org/0000-0001-5589-7896>

JORGE GARZA-VARGAS

Department of Mathematics, Princeton University, Princeton, NJ, USA

E-mail: jgarzavargas@princeton.edu

<https://orcid.org/0000-0001-6258-0600>

JOEL A. TROPP

Department of Computing and Mathematical Sciences, California Institute of Technol-
ogy, Pasadena, CA, USA

E-mail: jtropp@caltech.edu

<https://orcid.org/0000-0003-1024-1791>

RAMON VAN HANDEL

Department of Mathematics, Princeton University, Princeton, NJ, USA

E-mail: rvan@math.princeton.edu

<https://orcid.org/0000-0003-4026-8100>